

履 歴 書

令和4年1月

氏名 松島 綱治 (まつしま こうじ)

現在の職業 東京理科大学生命医科学研究所 炎症・免疫難病制御部門 教授
専門 炎症・免疫学

学歴 昭和53年3月 金沢大学医学部卒業
昭和53年5月 医籍登録 (第239651号)
昭和57年3月 金沢大学大学院医学研究科修了 (分子免疫学) 医学博士

職歴 昭和57年-昭和58年
Visiting Fellow, National Institute of Dental Research,
National Institutes of Health, USA
昭和58年-昭和60年
Visiting Fellow, National Cancer Institute
昭和60年-昭和62年
Visiting Associate, National Cancer Institute
昭和62年-平成2年
Visiting Scientist, National Cancer Institute
(平成元年 Tenure)
平成2年-平成9年
金沢大学 がん研究所 薬理部 教授
平成8年-平成30年
東京大学 大学院医学系研究科 分子予防医学 教授
平成30年-現在
東京理科大学 生命医科学研究所 炎症・免疫難病制御部門 教授

主な所属学会

The American Association of Immunologists
International Cytokine and Interferon Society
International Association of Inflammation Societies (President, 2011-2013)
日本癌学会
日本免疫学会 (理事, 1999-2002, 2004-2008 and 2010-2014)
日本炎症・再生医学会 (理事, 2001-present. 理事長, 2010-2012)
マクロファージ分子細胞生物学会 (会長, 1995-2016)
日本インターフェロン・サイトカイン学会 (会長, 2006-2008)

受賞歴

- 1991-Public Health Special Recognition Award from DHHS, USA. Interleukin 8 and monocyte chemotactic and activating factor の発見、精製、遺伝子クローニング
- 1998-高岡市民文化賞
- 2012-日本がん学会 JCA-CHAAO 賞 (anti-CCR4Ab、共同受賞)
- 2015-日本薬学会創薬科学賞 (anti-CCR4Ab、共同受賞)
- 2016-文部科学大臣表彰 科学技術賞 開発部門 (anti-CCR4Ab、共同受賞)
- 2016-日本免疫学会ヒト免疫研究賞 (サイトカイン・ケモカインの基礎研究を通じた免疫難病治療への貢献)
- 2019-International Cytokine and Interferon Society The Lifetime Honorary Membership Award
- 2020-東レ科学技術賞 (炎症制御因子研究を通じた免疫難病治療への貢献)
- 2021-武田医学賞 (ケモカインの発見による白血球浸潤機序の解明と創薬への貢献)

原著論文ならびに英語での総説

1. Hirose Y, Konda S, Sugai S, Takiguchi T, Shimizu S, Matsushima K, Annen Y. A case of eosinophilic leukemia associated with Loffler's endocarditis. Clinical course and autopsy findings. *Nippon Ketsueki Gakkai Zasshi*. 44:902-909, 1981.
2. Matsushima K, Cheng M, Migita S. Purification and physicochemical characterization of human alpha 2-HS-glycoprotein. *Biochim Biophys Acta*. 701:200-205, 1982.
3. Akiyama Y, Stevenson GW, Schlick E, Matsushima K, Miller PJ, Stevenson HC. Differential ability of human blood monocyte subsets to release various cytokines. *J Leukoc Biol*. 37:519-530, 1985.
4. Auron PE, Rosenwasser LJ, Matsushima K, Copeland T, Dinarello CA, Oppenheim JJ, Webb AC. Human and murine interleukin 1 possess sequence and structural similarities. *J Mol Cell Immunol*. 2:169-177, 1985.
5. Matsushima K, Bano M, Kidwell WR, Oppenheim JJ. Interleukin 1 increases collagen type IV production by murine mammary epithelial cells. *J Immunol*. 134:904-909, 1985.
6. Matsushima K, Durum SK, Kimball ES, Oppenheim JJ. Purification of human interleukin 1 from human monocyte culture supernatants and identity of thymocyte comitogenic factor, fibroblast-proliferation factor, acute-phase protein-inducing factor, and endogenous pyrogen. *Cell Immunol*. 92:290-301, 1985.
7. Matsushima K, Kuang YD, Tosato G, Hopkins SJ, Oppenheim JJ. B-cell-derived interleukin 1 (IL-1)-like factor. I. Relationship of production of IL-1-like factor to accessory cell function of Epstein-Barr virus-transformed human B-lymphoblast lines. *Cell Immunol*. 94:406-417, 1985.
8. Matsushima K, Oppenheim JJ. Calcium ionophore (A23187) increases interleukin 1 (IL-1) production by human peripheral blood monocytes and interacts synergistically with IL-1 to augment concanavalin A stimulated thymocyte proliferation. *Cell Immunol*. 90:226-233, 1985.
9. Matsushima K, Procopio A, Abe H, Scala G, Ortaldo JR, Oppenheim JJ. Production of interleukin 1 activity by normal human peripheral blood B lymphocytes. *J Immunol*.

135:1132-1136, 1985.

10. Matsushima K, Tosato G, Benjamin D, Oppenheim JJ. B-cell-derived interleukin-1 (IL-1)-like factor. II. Sources, effects, and biochemical properties. *Cell Immunol.* 94:418-426, 1985.
11. Onozaki K, Matsushima K, Aggarwal BB, Oppenheim JJ. Human interleukin 1 is a cytotoxic factor for several tumor cell lines. *J Immunol.* 135:3962-3968, 1985.
12. Onozaki K, Matsushima K, Kleinerman ES, Saito T, Oppenheim JJ. Role of interleukin 1 in promoting human monocyte-mediated tumor cytotoxicity. *J Immunol.* 135:314-320, 1985.
13. Abe H, Rossio JL, Ruscetti FW, Matsushima K, Oppenheim JJ. Establishment of a human B cell line that proliferates in response to B cell growth factor. *J Immunol Methods.* 90:111-123, 1986.
14. Endo Y, Matsushima K, Oppenheim JJ. Mechanism of in vitro antitumor effects of interleukin 1 (IL 1). *Immunobiology.* 172:316-322, 1986.
15. Matsushima K, Akahoshi T, Yamada M, Furutani Y, Oppenheim JJ. Properties of a specific interleukin 1 (IL 1) receptor on human Epstein Barr virus-transformed B lymphocytes: identity of the receptor for IL 1-alpha and IL 1-beta. *J Immunol.* 136:4496-4502, 1986.
16. Matsushima K, Copeland TD, Onozaki K, Oppenheim JJ. Purification and biochemical characteristics of two distinct human interleukins 1 from the myelomonocytic THP-1 cell line. *Biochemistry.* 25:3424-3429, 1986.
17. Matsushima K, Taguchi M, Kovacs EJ, Young HA, Oppenheim JJ. Intracellular localization of human monocyte associated interleukin 1 (IL 1) activity and release of biologically active IL 1 from monocytes by trypsin and plasmin. *J Immunol.* 136:2883-2891, 1986.
18. Matsushima K, Yodoi J, Tagaya Y, Oppenheim JJ. Down-regulation of interleukin 1 (IL 1) receptor expression by IL 1 and fate of internalized ¹²⁵I-labeled IL 1 beta in a human large granular lymphocyte cell line. *J Immunol.* 137:3183-3188, 1986.
19. Oppenheim JJ, Kovacs EJ, Matsushima K, Durum SK. There is more than one interleukin 1. *Immunol. Today* 2: 45-56, 1986.
20. Kasahara T, Mukaida N, Shinomiya H, Imai M, Matsushima K, Wakasugi H, Nakano K. Preparation and characterization of polyclonal and monoclonal antibodies against human interleukin 1 alpha (IL 1 alpha). *J Immunol.* 138:1804-1812, 1987.
21. Kawano M, Matsushima K, Oppenheim JJ. Identification of a major 50-kDa molecular weight human B-cell growth factor with Tac antigen-inducing activity on B cells. *Cell Immunol.* 108:132-149, 1987.
22. Matsushima K, Akahoshi T, Oppenheim JJ. Regulation of interleukin-1 (IL-1) receptor expression and protein phosphorylation induced by IL-1 stimulation. *Ann Inst Pasteur Immunol.* 138:478-481, 1987.
23. Matsushima K, Kobayashi Y, Copeland TD, Akahoshi T, Oppenheim JJ. Phosphorylation of a cytosolic 65-kDa protein induced by interleukin 1 in glucocorticoid pretreated normal human peripheral blood mononuclear leukocytes. *J Immunol.* 139:3367-3374, 1987.
24. Onozaki K, Tamatani T, Hashimoto T, Matsushima K. Growth inhibition and augmentation of mouse myeloid leukemic cell line differentiation by interleukin 1. *Cancer Res.*

- 47:2397-2402, 1987.
25. Oppenheim JJ, Matsushima K, Yoshimura T, Leonard EJ. The activities of cytokines are pleiotropic and interdependent. *Immunol Lett.* 16:179-183, 1987.
 26. Yodoi J, Okada M, Tagaya Y, Taniguchi Y, Teshigawara K, Kasahara T, Dinarello CA, Matsushima K, Honko T, Uchiyama T, et al. IL-2 receptor gene activation by ATL-derived factor (ADF). *Adv Exp Med Biol.* 213:139-148, 1987.
 27. Yoshimura T, Matsushima K, Oppenheim JJ, Leonard EJ. Neutrophil chemotactic factor produced by lipopolysaccharide (LPS)-stimulated human blood mononuclear leukocytes: partial characterization and separation from interleukin 1 (IL 1). *J Immunol.* 139:788-793, 1987.
 28. Yoshimura T, Matsushima K, Tanaka S, Robinson EA, Appella E, Oppenheim JJ, Leonard EJ. Purification of a human monocyte-derived neutrophil chemotactic factor that has peptide sequence similarity to other host defense cytokines. *Proc Natl Acad Sci U S A.* 84:9233-9237, 1987.
 29. Akahoshi T, Oppenheim JJ, Matsushima K. Interleukin 1 stimulates its own receptor expression on human fibroblasts through the endogenous production of prostaglandin(s). *J Clin Invest.* 82:1219-1224, 1988.
 30. Akahoshi T, Oppenheim JJ, Matsushima K. Induction of high-affinity interleukin 1 receptor on human peripheral blood lymphocytes by glucocorticoid hormones. *J Exp Med.* 167:924-936, 1988.
 31. Endo Y, Matsushima K, Onozaki K, Oppenheim JJ. Role of ornithine decarboxylase in the regulation of cell growth by IL-1 and tumor necrosis factor. *J Immunol.* 141:2342-2348, 1988.
 32. Kawano M, Matsushima K, Masuda A, Oppenheim JJ. A major 50-kDa human B-cell growth factor-II induces both Tac antigen expression and proliferation by several types of lymphocytes. *Cell Immunol.* 111:273-286, 1988.
 33. Kobayashi Y, Appella E, Yamada M, Copeland TD, Oppenheim JJ, Matsushima K. Phosphorylation of intracellular precursors of human IL-1. *J Immunol.* 140:2279-2287, 1988.
 34. Lavu S, Clark J, Swarup R, Matsushima K, Paturu K, Moss J, Kung HF. Molecular cloning and DNA sequence analysis of the human guanine nucleotide-binding protein Go alpha. *Biochem Biophys Res Commun.* 150:811-815, 1988.
 35. Lew W, Oppenheim JJ, Matsushima K. Analysis of the suppression of IL-1 alpha and IL-1 beta production in human peripheral blood mononuclear adherent cells by a glucocorticoid hormone. *J Immunol.* 140:1895-1902, 1988.
 36. Masuda A, Longo DL, Kobayashi Y, Appella E, Oppenheim JJ, Matsushima K. Induction of mitochondrial manganese superoxide dismutase by interleukin 1. *Faseb J.* 2:3087-3091, 1988.
 37. Matsushima K, Morishita K, Yoshimura T, Lavu S, Kobayashi Y, Lew W, Appella E, Kung HF, Leonard EJ, Oppenheim JJ. Molecular cloning of a human monocyte-derived neutrophil chemotactic factor (MDNCF) and the induction of MDNCF mRNA by interleukin 1 and tumor necrosis factor. *J Exp Med.* 167:1883-1893, 1988.

38. Matsushima K, Shiroo M, Kung HF, Copeland TD. Purification and characterization of a cytosolic 65-kilodalton phosphoprotein in human leukocytes whose phosphorylation is augmented by stimulation with interleukin 1. *Biochemistry*. 27:3765-3770, 1988.
39. Merrill JE, Matsushima K. Production of and response to interleukin 1 by cloned human oligodendrogloma cell lines. *J Biol Regul Homeost Agents*. 2:77-86, 1988.
40. Modi WS, Masuda A, Yamada M, Oppenheim JJ, Matsushima K, O'Brien SJ. Chromosomal localization of the human interleukin 1 alpha (IL-1 alpha) gene. *Genomics*. 2:310-314, 1988.
41. Oppenheim JJ, Lew W, Akahoshi T, Matsushima K, Neta R. Aspects of cytokine induced modulation of immunity and inflammation with emphasis on interleukin 1. *Arzneimittelforschung*. 38:461-465, 1988.
42. Tagaya Y, Okada M, Sugie K, Kasahara T, Kondo N, Hamuro J, Matsushima K, Dinarello CA, Yodoi J. IL-2 receptor(p55)/Tac-inducing factor. Purification and characterization of adult T cell leukemia-derived factor. *J Immunol*. 140:2614-2620, 1988.
43. Tanaka S, Robinson EA, Yoshimura T, Matsushima K, Leonard EJ, Appella E. Synthesis and biological characterization of monocyte-derived neutrophil chemotactic factor. *FEBS Lett*. 236:467-470, 1988.
44. Clore GM, Appella E, Yamada M, Matsushima K, Gronenborn AM. Determination of the secondary structure of interleukin-8 by nuclear magnetic resonance spectroscopy. *J Biol Chem*. 264:18907-18911, 1989.
45. Furutani Y, Nomura H, Notake M, Oyamada Y, Fukui T, Yamada M, Larsen CG, Oppenheim JJ, Matsushima K. Cloning and sequencing of the cDNA for human monocyte chemotactic and activating factor (MCAF). *Biochem Biophys Res Commun*. 159:249-255, 1989.
46. Larsen CG, Anderson AO, Appella E, Oppenheim JJ, Matsushima K. The neutrophil-activating protein (NAP-1) is also chemotactic for T lymphocytes. *Science*. 243:1464-1466, 1989.
47. Larsen CG, Anderson AO, Oppenheim JJ, Matsushima K. Production of interleukin-8 by human dermal fibroblasts and keratinocytes in response to interleukin-1 or tumour necrosis factor. *Immunology*. 68:31-36, 1989.
48. Larsen CG, Zachariae CO, Oppenheim JJ, Matsushima K. Production of monocyte chemotactic and activating factor (MCAF) by human dermal fibroblasts in response to interleukin 1 or tumor necrosis factor. *Biochem Biophys Res Commun*. 160:1403-1408, 1989.
49. Matsushima K, Larsen CG, DuBois GC, Oppenheim JJ. Purification and characterization of a novel monocyte chemotactic and activating factor produced by a human myelomonocytic cell line. *J Exp Med*. 169:1485-1490, 1989.
50. Matsushima K, Oppenheim JJ. Interleukin 8 and MCAF: novel inflammatory cytokines inducible by IL 1 and TNF. *Cytokine*. 1:2-13, 1989.
51. Morinaga Y, Suzuki H, Takatsuki F, Akiyama Y, Taniyama T, Matsushima K, Onozaki K. Contribution of IL-6 to the antiproliferative effect of IL-1 and tumor necrosis factor on tumor cell lines. *J Immunol*. 143:3538-3542, 1989.
52. Mukaida N, Shiroo M, Matsushima K. Genomic structure of the human monocyte-derived

- neutrophil chemotactic factor IL-8. *J Immunol.* 143:1366-1371, 1989.
53. Nakao S, Matsushima K, Young N. Decreased interleukin 1 production in aplastic anaemia. *Br J Haematol.* 71:431-436, 1989.
 54. Oppenheim JJ, Matsushima K, Yoshimura T, Leonard EJ, Neta R. Relationship between interleukin 1 (IL1), tumor necrosis factor (TNF) and a neutrophil attracting peptide (NAP-1). *Agents Actions.* 26:134-140, 1989.
 55. Samanta AK, Oppenheim JJ, Matsushima K. Identification and characterization of specific receptors for monocyte-derived neutrophil chemotactic factor (MDNCF) on human neutrophils. *J Exp Med.* 169:1185-1189, 1989.
 56. Yoshimura T, Robinson EA, Appella E, Matsushima K, Showalter SD, Skeel A, Leonard EJ. Three forms of monocyte-derived neutrophil chemotactic factor (MDNCF) distinguished by different lengths of the amino-terminal sequence. *Mol Immunol.* 26:87-93, 1989.
 57. Appella E, Matsushima K, Oppenheim JJ, Yoshimura T, Leonard EJ, Clore GM, Gronenborn AM. Determination of the primary and secondary structure of NAP-1/IL-8 and a monocyte chemoattractant protein, MCP-1/MCAF. *Prog Clin Biol Res.* 349:405-417, 1990.
 58. Baldwin ET, Franklin KA, Appella E, Yamada M, Matsushima K, Wlodawer A, Weber IT. Crystallization of human interleukin-8. A protein chemotactic for neutrophils and T-lymphocytes. *J Biol Chem.* 265:6851-6853, 1990.
 59. Balentien E, Han JH, Thomas HG, Wen DZ, Samantha AK, Zachariae CO, Griffin PR, Brachmann R, Wong WL, Matsushima K, Richmond A. Recombinant expression, biochemical characterization, and biological activities of the human MGSA/gro protein. *Biochemistry.* 29:10225-10233, 1990.
 60. Brennan FM, Zachariae CO, Chantry D, Larsen CG, Turner M, Maini RN, Matsushima K, Feldmann M. Detection of interleukin 8 biological activity in synovial fluids from patients with rheumatoid arthritis and production of interleukin 8 mRNA by isolated synovial cells. *Eur J Immunol.* 20:2141-2144, 1990.
 61. Clore GM, Appella E, Yamada M, Matsushima K, Gronenborn AM. Three-dimensional structure of interleukin 8 in solution. *Biochemistry.* 29:1689-1696, 1990.
 62. Djeu JY, Matsushima K, Oppenheim JJ, Shiotsuki K, Blanchard DK. Functional activation of human neutrophils by recombinant monocyte-derived neutrophil chemotactic factor/IL-8. *J Immunol.* 144:2205-2210, 1990.
 63. Hirose K, Zachariae CO, Oppenheim JJ, Matsushima K. Induction of gene expression and production of immunomodulating cytokines by PSK in human peripheral blood mononuclear cells. *Lymphokine Res.* 9:475-483, 1990.
 64. Kobayashi Y, Oppenheim JJ, Matsushima K. Calcium-dependent binding of phosphorylated human pre interleukin 1 alpha to phospholipids. *J Biochem (Tokyo).* 107:666-670, 1990.
 65. Kobayashi Y, Yamamoto K, Saïdo T, Kawasaki H, Oppenheim JJ, Matsushima K. Identification of calcium-activated neutral protease as a processing enzyme of human interleukin 1 alpha. *Proc Natl Acad Sci U S A.* 87:5548-5552, 1990.
 66. Larsen C, Zachariae C, Mukaida N, Anderson A, Yamada M, Oppenheim J, Matsushima K. Proinflammatory cytokines interleukin 1 and tumor necrosis factor induce cytokines that are

- chemotactic for neutrophils, T cells and monocytes. *Prog Clin Biol Res.* 349:419-431, 1990.
67. Modi WS, Dean M, Seuanez HN, Mukaida N, Matsushima K, O'Brien SJ. Monocyte-derived neutrophil chemotactic factor (MDNCF/IL-8) resides in a gene cluster along with several other members of the platelet factor 4 gene superfamily. *Hum Genet.* 84:185-187, 1990.
 68. Mukaida N, Mahe Y, Matsushima K. Cooperative interaction of nuclear factor-kappa B- and cis-regulatory enhancer binding protein-like factor binding elements in activating the interleukin-8 gene by pro-inflammatory cytokines. *J Biol Chem.* 265:21128-21133, 1990.
 69. Samanta AK, Oppenheim JJ, Matsushima K. Interleukin 8 (monocyte-derived neutrophil chemotactic factor) dynamically regulates its own receptor expression on human neutrophils. *J Biol Chem.* 265:183-189, 1990.
 70. Shiroo M, Matsushima K. Enhanced phosphorylation of 65 and 74 kDa proteins by tumor necrosis factor and interleukin-1 in human peripheral blood mononuclear cells. *Cytokine.* 2:13-20, 1990.
 71. Sica A, Matsushima K, Van Damme J, Wang JM, Polentarutti N, Dejana E, Colotta F, Mantovani A. IL-1 transcriptionally activates the neutrophil chemotactic factor/IL-8 gene in endothelial cells. *Immunology.* 69:548-553, 1990.
 72. Sica A, Wang JM, Colotta F, Dejana E, Mantovani A, Oppenheim JJ, Larsen CG, Zachariae CO, Matsushima K. Monocyte chemotactic and activating factor gene expression induced in endothelial cells by IL-1 and tumor necrosis factor. *J Immunol.* 144:3034-3038, 1990.
 73. Wang JM, Taraboletti G, Matsushima K, Van Damme J, Mantovani A. Induction of haptotactic migration of melanoma cells by neutrophil activating protein/interleukin-8. *Biochem Biophys Res Commun.* 169:165-170, 1990.
 74. Zachariae CO, Anderson AO, Thompson HL, Appella E, Mantovani A, Oppenheim JJ, Matsushima K. Properties of monocyte chemotactic and activating factor (MCAF) purified from a human fibrosarcoma cell line. *J Exp Med.* 171:2177-2182, 1990.
 75. Abbott F, Ryan JJ, Ceska M, Matsushima K, Sarraf CE, Rees AJ. Interleukin-1 beta stimulates human mesangial cells to synthesize and release interleukins-6 and -8. *Kidney Int.* 40:597-605, 1991.
 76. Baldwin ET, Weber IT, St Charles R, Xuan JC, Appella E, Yamada M, Matsushima K, Edwards BF, Clore GM, Gronenborn AM, et al. Crystal structure of interleukin 8: symbiosis of NMR and crystallography. *Proc Natl Acad Sci U S A.* 88:502-506, 1991.
 77. Endo H, Akahoshi T, Takagishi K, Kashiwazaki S, Matsushima K. Elevation of interleukin-8 (IL-8) levels in joint fluids of patients with rheumatoid arthritis and the induction by IL-8 of leukocyte infiltration and synovitis in rabbit joints. *Lymphokine Cytokine Res.* 10:245-252, 1991.
 78. Ferrick MR, Thureau SR, Oppenheim MH, Herbort CP, Ni M, Zachariae CO, Matsushima K, Chan CC. Ocular inflammation stimulated by intravitreal interleukin-8 and interleukin-1. *Invest Ophthalmol Vis Sci.* 32:1534-1539, 1991.
 79. Kasahara T, Mukaida N, Yamashita K, Yagisawa H, Akahoshi T, Matsushima K. IL-1 and TNF-alpha induction of IL-8 and monocyte chemotactic and activating factor (MCAF) mRNA expression in a human astrocytoma cell line. *Immunology.* 74:60-67, 1991.

80. Kobayashi Y, Oppenheim JJ, Matsushima K. Human pre-interleukin 1 alpha and beta: structural features revealed by limited proteolysis. *Chem Pharm Bull (Tokyo)*. 39:1513-1517, 1991.
81. Kudo C, Araki A, Matsushima K, Sendo F. Inhibition of IL-8-induced W3/25+ (CD4+) T lymphocyte recruitment into subcutaneous tissues of rats by selective depletion of in vivo neutrophils with a monoclonal antibody. *J Immunol*. 147:2196-2201, 1991.
82. Kuno K, Kuno S, Matsushima K, Murakami S. Evidence for binding of at least two factors, including T-rich strand-binding factor(s) to the single-stranded ARS1 sequence in *Saccharomyces cerevisiae*. *Mol Gen Genet*. 230:45-48, 1991.
83. Larsen CG, Kristensen M, Paludan K, Deleuran B, Thomsen MK, Zachariae C, Kragballe K, Matsushima K, Thestrup-Pedersen K. 1,25(OH)₂-D₃ is a potent regulator of interleukin-1 induced interleukin-8 expression and production. *Biochem Biophys Res Commun*. 176:1020-1026, 1991.
84. Mahe Y, Mukaida N, Kuno K, Akiyama M, Ikeda N, Matsushima K, Murakami S. Hepatitis B virus X protein transactivates human interleukin-8 gene through acting on nuclear factor kB and CCAAT/enhancer-binding protein-like cis-elements. *J Biol Chem*. 266:13759-13763, 1991.
85. Matsushime H, Roussel MF, Matsushima K, Hishinuma A, Sherr CJ. Cloning and expression of murine interleukin-1 receptor antagonist in macrophages stimulated by colony-stimulating factor 1. *Blood*. 78:616-623, 1991.
86. Mukaida N, Hishinuma A, Zachariae CO, Oppenheim JJ, Matsushima K. Regulation of human interleukin 8 gene expression and binding of several other members of the intercrine family to receptors for interleukin-8. *Adv Exp Med Biol*. 305:31-38, 1991.
87. Mukaida N, Zachariae CC, Gusella GL, Matsushima K. Dexamethasone inhibits the induction of monocyte chemotactic-activating factor production by IL-1 or tumor necrosis factor. *J Immunol*. 146:1212-1215, 1991.
88. Oppenheim JJ, Zachariae CO, Mukaida N, Matsushima K. Properties of the novel proinflammatory supergene "intercrine" cytokine family. *Annu Rev Immunol*. 9:617-648, 1991.
89. Smyth MJ, Zachariae CO, Norihisa Y, Ortaldo JR, Hishinuma A, Matsushima K. IL-8 gene expression and production in human peripheral blood lymphocyte subsets. *J Immunol*. 146:3815-3823, 1991.
90. Sozzani S, Luini W, Molino M, Jilek P, Bottazzi B, Cerletti C, Matsushima K, Mantovani A. The signal transduction pathway involved in the migration induced by a monocyte chemotactic cytokine. *J Immunol*. 147:2215-2221, 1991.
91. Terkeltaub R, Zachariae C, Santoro D, Martin J, Peveri P, Matsushima K. Monocyte-derived neutrophil chemotactic factor/interleukin-8 is a potential mediator of crystal-induced inflammation. *Arthritis Rheum*. 34:894-903, 1991.
92. Tsai WP, Hirose K, Nara PL, Kuang YD, Conley S, Li BQ, Kung HF, Matsushima K. Decrease in cytokine production by HIV-infected macrophages in response to LPS-mediated activation. *Lymphokine Cytokine Res*. 10:421-429, 1991.

93. Yuo A, Kitagawa S, Kasahara T, Matsushima K, Saito M, Takaku F. Stimulation and priming of human neutrophils by interleukin-8: cooperation with tumor necrosis factor and colony-stimulating factors. *Blood*. 78:2708-2714, 1991.
94. Zachariae CO, Thestrup-Pedersen K, Matsushima K. Expression and secretion of leukocyte chemotactic cytokines by normal human melanocytes and melanoma cells. *J Invest Dermatol*. 97:593-599, 1991.
95. Ko Y, Mukaida N, Panyutich A, Voitenok NN, Matsushima K, Kawai T, Kasahara T. A sensitive enzyme-linked immunosorbent assay for human interleukin-8. *J Immunol Methods*. 149:227-235, 1992.
96. Matsushima K, Baldwin ET, Mukaida N. Interleukin-8 and MCAF: novel leukocyte recruitment and activating cytokines. *Chem Immunol*. 51:236-265, 1992.
97. Mukaida N, Gussella GL, Kasahara T, Ko Y, Zachariae CO, Kawai T, Matsushima K. Molecular analysis of the inhibition of interleukin-8 production by dexamethasone in a human fibrosarcoma cell line. *Immunology*. 75:674-679, 1992.
98. Mukaida N, Harada A, Yasumoto K, Matsushima K. Properties of pro-inflammatory cell type-specific leukocyte chemotactic cytokines, interleukin 8 (IL-8) and monocyte chemotactic and activating factor (MCAF). *Microbiol Immunol*. 36:773-789, 1992.
99. Mukaida N, Matsushima K. Regulation of IL-8 production and the characteristics of the receptors for IL-8. *Cytokines*. 4:41-53, 1992.
100. Nourshargh S, Perkins JA, Showell HJ, Matsushima K, Williams TJ, Collins PD. A comparative study of the neutrophil stimulatory activity in vitro and pro-inflammatory properties in vivo of 72 amino acid and 77 amino acid IL-8. *J Immunol*. 148:106-111, 1992.
101. Sakamoto K, Masuda T, Mita S, Ishiko T, Nakashima Y, Arakawa H, Egami H, Harada S, Matsushima K, Ogawa M. Interleukin-8 is constitutively and commonly produced by various human carcinoma cell lines. *Int J Clin Lab Res*. 22:216-219, 1992.
102. Thompson HL, Matsushima K. Human polymorphonuclear leucocytes stimulated by tumour necrosis factor-alpha show increased adherence to extracellular matrix proteins which is mediated via the CD11b/18 complex. *Clin Exp Immunol*. 90:280-285, 1992.
103. Yasumoto K, Okamoto S, Mukaida N, Murakami S, Mai M, Matsushima K. Tumor necrosis factor alpha and interferon gamma synergistically induce interleukin 8 production in a human gastric cancer cell line through acting concurrently on AP-1 and NF-kB-like binding sites of the interleukin 8 gene. *J Biol Chem*. 267:22506-22511, 1992.
104. Abe Y, Kawakami M, Kuroki M, Yamamoto T, Fujii M, Kobayashi H, Yaginuma T, Kashii A, Saito M, Matsushima K. Transient rise in serum interleukin-8 concentration during acute myocardial infarction. *Br Heart J*. 70:132-134, 1993.
105. Akahoshi T, Wada C, Endo H, Hirota K, Hosaka S, Takagishi K, Kondo H, Kashiwazaki S, Matsushima K. Expression of monocyte chemotactic and activating factor in rheumatoid arthritis. Regulation of its production in synovial cells by interleukin-1 and tumor necrosis factor. *Arthritis Rheum*. 36:762-771, 1993.
106. Asano T, Fujimaki W, McWatters A, An T, Matsushima K, Kleinerman ES. Effect of Adriamycin on liposomal muramyl tripeptide's ability to up-regulate monocyte cytokine

- expression. *Cancer Immunol Immunother.* 37:408-411, 1993.
107. Gusella GL, Musso T, Bosco MC, Espinoza-Delgado I, Matsushima K, Varesio L. IL-2 up-regulates but IFN-gamma suppresses IL-8 expression in human monocytes. *J Immunol.* 151:2725-2732, 1993.
 108. Harada A, Sekido N, Kuno K, Akiyama M, Kasahara T, Nakanishi I, Mukaida N, Matsushima K. Expression of recombinant rabbit IL-8 in *Escherichia coli* and establishment of the essential involvement of IL-8 in recruiting neutrophils into lipopolysaccharide-induced inflammatory site of rabbit skin. *Int Immunol.* 5:681-690, 1993.
 109. Hirose K, Longo DL, Oppenheim JJ, Matsushima K. Overexpression of mitochondrial manganese superoxide dismutase promotes the survival of tumor cells exposed to interleukin-1, tumor necrosis factor, selected anticancer drugs, and ionizing radiation. *Faseb J.* 7:361-368, 1993.
 110. Horiguchi H, Mukaida N, Okamoto S, Teranishi H, Kasuya M, Matsushima K. Cadmium induces interleukin-8 production in human peripheral blood mononuclear cells with the concomitant generation of superoxide radicals. *Lymphokine Cytokine Res.* 12:421-428, 1993.
 111. Jinquan T, Larsen CG, Gesser B, Matsushima K, Thestrup-Pedersen K. Human IL-10 is a chemoattractant for CD8+ T lymphocytes and an inhibitor of IL-8-induced CD4+ T lymphocyte migration. *J Immunol.* 151:4545-4551, 1993.
 112. Ko YC, Mukaida N, Ishiyama S, Tokue A, Kawai T, Matsushima K, Kasahara T. Elevated interleukin-8 levels in the urine of patients with urinary tract infections. *Infect Immun.* 61:1307-1314, 1993.
 113. Kristensen M, Jinquan T, Thomsen MK, Zachariae C, Paludan K, Ahnfelt-Ronne I, Matsushima K, Thestrup-Pedersen K, Larsen CG. ETH615, a synthetic inhibitor of leukotriene biosynthesis and function, also inhibits the production of and biological responses towards interleukin-8. *Exp Dermatol.* 2:165-170, 1993.
 114. Kuno K, Okamoto S, Hirose K, Murakami S, Matsushima K. Structure and function of the intracellular portion of the mouse interleukin 1 receptor (type I). Determining the essential region for transducing signals to activate the interleukin 8 gene. *J Biol Chem.* 268:13510-13518, 1993.
 115. Matsusaka T, Fujikawa K, Nishio Y, Mukaida N, Matsushima K, Kishimoto T, Akira S. Transcription factors NF-IL6 and NF-kappa B synergistically activate transcription of the inflammatory cytokines, interleukin 6 and interleukin 8. *Proc Natl Acad Sci U S A.* 90:10193-10197, 1993.
 116. Matsushima K, Sone S, Onozaki K, Yamazaki M, Yoshida T, Irimura T. Meeting report of the International Symposium of Molecular Cell Biology of Macrophages '92. *Cytokine.* 5:91-94, 1993.
 117. Morita M, Kasahara T, Mukaida N, Matsushima K, Nagashima T, Nishizawa M, Yoshida M. Induction and regulation of IL-8 and MCAF production in human brain tumor cell lines and brain tumor tissues. *Eur Cytokine Netw.* 4:351-358, 1993.
 118. Nomura H, Nielsen BW, Matsushima K. Molecular cloning of cDNAs encoding a LD78

- receptor and putative leukocyte chemotactic peptide receptors. *Int Immunol.* 5:1239-1249, 1993.
119. Okamoto S, Mukaida N, Yasumoto K, Horiguchi H, Matsushima K. Molecular mechanism of interleukin-8 gene expression. *Adv Exp Med Biol.* 351:87-97, 1993.
 120. Sekido N, Mukaida N, Harada A, Nakanishi I, Watanabe Y, Matsushima K. Prevention of lung reperfusion injury in rabbits by a monoclonal antibody against interleukin-8. *Nature.* 365:654-657, 1993.
 121. Singh RK, Berry K, Matsushima K, Yasumoto K, Fidler IJ. Synergism between human monocyte chemotactic and activating factor and bacterial products for activation of tumoricidal properties in murine macrophages. *J Immunol.* 151:2786-2793, 1993.
 122. Taub DD, Lloyd AR, Conlon K, Wang JM, Ortaldo JR, Harada A, Matsushima K, Kelvin DJ, Oppenheim JJ. Recombinant human interferon-inducible protein 10 is a chemoattractant for human monocytes and T lymphocytes and promotes T cell adhesion to endothelial cells. *J Exp Med.* 177:1809-1814, 1993.
 123. Tsuchiya I, Kasahara T, Yamashita K, Ko YC, Kanazawa K, Matsushima K, Mukaida N. Induction of inflammatory cytokines in the pleural effusion of cancer patients after the administration of an immunomodulator, OK-432: role of IL-8 for neutrophil infiltration. *Cytokine.* 5:595-603, 1993.
 124. Wang JM, Hishinuma A, Oppenheim JJ, Matsushima K. Studies of binding and internalization of human recombinant monocyte chemotactic and activating factor (MCAF) by monocytic cells. *Cytokine.* 5:264-275, 1993.
 125. Akahoshi T, Endo H, Kondo H, Kashiwazaki S, Kasahara T, Mukaida N, Harada A, Matsushima K. Essential involvement of interleukin-8 in neutrophil recruitment in rabbits with acute experimental arthritis induced by lipopolysaccharide and interleukin-1. *Lymphokine Cytokine Res.* 13:113-116, 1994.
 126. Asano T, Matsushima K, Kleinerman ES. Liposome-encapsulated muramyl tripeptide up-regulates monocyte chemotactic and activating factor gene expression in human monocytes at the transcriptional and post-transcriptional levels. *Cancer Immunol Immunother.* 38:16-22, 1994.
 127. Asano T, McWatters A, An T, Matsushima K, Kleinerman ES. Liposomal muramyl tripeptide up-regulates interleukin-1 alpha, interleukin-1 beta, tumor necrosis factor-alpha, interleukin-6 and interleukin-8 gene expression in human monocytes. *J Pharmacol Exp Ther.* 268:1032-1039, 1994.
 128. Deleuran B, Lemche P, Kristensen M, Chu CQ, Field M, Jensen J, Matsushima K, Stengaard-Pedersen K. Localisation of interleukin 8 in the synovial membrane, cartilage-pannus junction and chondrocytes in rheumatoid arthritis. *Scand J Rheumatol.* 23:2-7, 1994.
 129. Endo H, Akahoshi T, Nishimura A, Tonegawa M, Takagishi K, Kashiwazaki S, Matsushima K, Kondo H. Experimental arthritis induced by continuous infusion of IL-8 into rabbit knee joints. *Clin Exp Immunol.* 96:31-35, 1994.
 130. Garin EH, Blanchard DK, Matsushima K, Djeu JY. IL-8 production by peripheral blood

- mononuclear cells in nephrotic patients. *Kidney Int.* 45:1311-1317, 1994.
131. Harada A, Kuno K, Nomura H, Mukaida N, Murakami S, Matsushima K. Cloning of a cDNA encoding a mouse homolog of the interleukin-8 receptor. *Gene.* 142:297-300, 1994.
 132. Harada A, Sekido N, Akahoshi T, Wada T, Mukaida N, Matsushima K. Essential involvement of interleukin-8 (IL-8) in acute inflammation. *J Leukoc Biol.* 56:559-564, 1994.
 133. Ito A, Imada K, Sato T, Kubo T, Matsushima K, Mori Y. Suppression of interleukin 8 production by progesterone in rabbit uterine cervix. *Biochem J.* 301 (Pt 1):183-186, 1994.
 134. Itoh R, Takenaka T, Okitsu-Negishi S, Matsushima K, Mizoguchi M. Interleukin 8 in Behcet's disease. *J Dermatol.* 21:397-404, 1994.
 135. Kasahara K, Tobe T, Tomita M, Mukaida N, Shao-Bo S, Matsushima K, Yoshida T, Sugihara S, Kobayashi K. Selective expression of monocyte chemotactic and activating factor/monocyte chemoattractant protein 1 in human blood monocytes by *Mycobacterium tuberculosis*. *J Infect Dis.* 170:1238-1247, 1994.
 136. Kuno K, Matsushima K. The IL-1 receptor signaling pathway. *J Leukoc Biol.* 56:542-547, 1994.
 137. Kuno K, Sukegawa K, Ishikawa Y, Orii T, Matsushima K. Acid sphingomyelinase is not essential for the IL-1 and tumor necrosis factor receptor signaling pathway leading to NFkB activation. *Int Immunol.* 6:1269-1272, 1994.
 138. Matsushima K, Sone S, Onozaki K, Yamazaki M, Irimura T. Meeting report of the International Symposium of Molecular Cell Biology of Macrophages '94. *Lymphokine Cytokine Res.* 13:331-333, 1994.
 139. Mizuno K, Sone S, Orino E, Mukaida N, Matsushima K, Ogura T. Spontaneous production of interleukin-8 by human lung cancer cells and its augmentation by tumor necrosis factor alpha and interleukin-1 at protein and mRNA levels. *Oncology.* 51:467-471, 1994.
 140. Mukaida N, Morita M, Ishikawa Y, Rice N, Okamoto S, Kasahara T, Matsushima K. Novel mechanism of glucocorticoid-mediated gene repression. Nuclear factor-kappa B is target for glucocorticoid-mediated interleukin 8 gene repression. *J Biol Chem.* 269:13289-13295, 1994.
 141. Mukaida N, Okamoto S, Ishikawa Y, Matsushima K. Molecular mechanism of interleukin-8 gene expression. *J Leukoc Biol.* 56:554-558, 1994.
 142. Murakami S, Cheong J, Ohno S, Matsushima K, Kaneko S. Transactivation of human hepatitis B virus X protein, HBx, operates through a mechanism distinct from protein kinase C and okadaic acid activation pathways. *Virology.* 199:243-246, 1994.
 143. Murayama T, Kuno K, Jisaki F, Obuchi M, Sakamuro D, Furukawa T, Mukaida N, Matsushima K. Enhancement human cytomegalovirus replication in a human lung fibroblast cell line by interleukin-8. *J Virol.* 68:7582-7585, 1994.
 144. Murayama T, Tsuchiya N, Jisaki F, Ozaki M, Sakamuro D, Hirai K, Shimizu S, Ito K, Matsushima K, Furukawa T. Elevated cytokine levels in synovial fluid of rheumatoid arthritis correlates with the presence of cytomegalovirus genome. *Autoimmunity.* 17:333-337, 1994.
 145. Nakamoto Y, Kaneko S, Honda M, Unoura M, Cheong J, Harada A, Matsushima K,

- Kobayashi K, Murakami S. Detection of the putative E2 protein of hepatitis C virus in human liver. *J Med Virol.* 42:374-379, 1994.
146. Nakano Y, Kasahara T, Mukaida N, Ko YC, Nakano M, Matsushima K. Protection against lethal bacterial infection in mice by monocyte-chemotactic and -activating factor. *Infect Immun.* 62:377-383, 1994.
 147. Nielsen BW, Mukaida N, Matsushima K, Kasahara T. Macrophages as producers of chemotactic proinflammatory cytokines. *Immunol Ser.* 60:131-142, 1994.
 148. Oishi K, Sonoda F, Kobayashi S, Iwagaki A, Nagatake T, Matsushima K, Matsumoto K. Role of interleukin-8 (IL-8) and an inhibitory effect of erythromycin on IL-8 release in the airways of patients with chronic airway diseases. *Infect Immun.* 62:4145-4152, 1994.
 149. Okamoto S, Mukaida N, Yasumoto K, Rice N, Ishikawa Y, Horiguchi H, Murakami S, Matsushima K. The interleukin-8 AP-1 and kappa B-like sites are genetic end targets of FK506-sensitive pathway accompanied by calcium mobilization. *J Biol Chem.* 269:8582-8589, 1994.
 150. Oliveira IC, Mukaida N, Matsushima K, Vilcek J. Transcriptional inhibition of the interleukin-8 gene by interferon is mediated by the NF-kappa B site. *Mol Cell Biol.* 14:5300-5308, 1994.
 151. Wada T, Tomosugi N, Kobayashi K, Ishikawa I, Shinoda A, Tsugawa R, Mukaida N, Matsushima K. Appearance of interleukin-8 in the urine predicts acute rejection of renal transplantation. *Nephrol Dial Transplant.* 9:874-875, 1994.
 152. Wada T, Tomosugi N, Naito T, Yokoyama H, Kobayashi K, Harada A, Mukaida N, Matsushima K. Prevention of proteinuria by the administration of anti-interleukin 8 antibody in experimental acute immune complex-induced glomerulonephritis. *J Exp Med.* 180:1135-1140, 1994.
 153. Wada T, Yokoyama H, Tomosugi N, Hisada Y, Ohta S, Naito T, Kobayashi K, Mukaida N, Matsushima K. Detection of urinary interleukin-8 in glomerular diseases. *Kidney Int.* 46:455-460, 1994.
 154. Fujioka N, Mukaida N, Harada A, Akiyama M, Kasahara T, Kuno K, Ooi A, Mai M, Matsushima K. Preparation of specific antibodies against murine IL-1ra and the establishment of IL-1ra as an endogenous regulator of bacteria-induced fulminant hepatitis in mice. *J Leukoc Biol.* 58:90-98, 1995.
 155. Garin EH, West L, Blanchard K, Matsushima K, Djeu JY. Effect of lymphokines on 35sulfate uptake by the glomerular basement membrane. *Nephron.* 71:442-447, 1995.
 156. Hirose K, Hakozaiki M, Nyunoya Y, Kobayashi Y, Matsushita K, Takenouchi T, Mikata A, Mukaida N, Matsushima K. Chemokine gene transfection into tumour cells reduced tumorigenicity in nude mice in association with neutrophilic infiltration. *Br J Cancer.* 72:708-714, 1995.
 157. Hoshino Y, Hatake K, Kasahara T, Takahashi Y, Ikeda M, Tomizuka H, Ohtsuki T, Uwai M, Mukaida N, Matsushima K, et al. Monocyte chemoattractant protein-1 stimulates tumor necrosis and recruitment of macrophages into tumors in tumor-bearing nude mice: increased granulocyte and macrophage progenitors in murine bone marrow. *Exp Hematol.*

- 23:1035-1039, 1995.
158. Ishikawa Y, Mukaida N, Kuno K, Rice N, Okamoto S, Matsushima K. Establishment of lipopolysaccharide-dependent nuclear factor kappa B activation in a cell-free system. *J Biol Chem.* 270:4158-4164, 1995.
 159. Jinquan T, Frydenberg J, Mukaida N, Bonde J, Larsen CG, Matsushima K, Thestrup-Pedersen K. Recombinant human growth-regulated oncogene-alpha induces T lymphocyte chemotaxis. A process regulated via IL-8 receptors by IFN-gamma, TNF-alpha, IL-4, IL-10, and IL-13. *J Immunol.* 155:5359-5368, 1995.
 160. Ko YC, Mukaida N, Kasahara T, Muto S, Matsushima K, Kusano E, Asano Y, Itoh Y, Yamagishi Y, Kawai T. Specific increase in interleukin-8 concentrations in dialysis fluid of patients with peritonitis receiving continuous ambulatory peritoneal dialysis. *J Clin Pathol.* 48:115-119, 1995.
 161. Kuno K, Ishikawa Y, Ernst MK, Ogata M, Rice NR, Mukaida N, Matsushima K. Identification of an I kappa B alpha-associated protein kinase in a human monocytic cell line and determination of its phosphorylation sites on I kappa B alpha. *J Biol Chem.* 270:27914-27919, 1995.
 162. Morohashi H, Miyawaki T, Nomura H, Kuno K, Murakami S, Matsushima K, Mukaida N. Expression of both types of human interleukin-8 receptors on mature neutrophils, monocytes, and natural killer cells. *J Leukoc Biol.* 57:180-187, 1995.
 163. Mukaida N, Harada A, Matsushima K. A novel leukocyte chemotactic and activating cytokine, interleukin-8 (IL-8). *Cancer Treat Res.* 80:261-286, 1995.
 164. Nakashima E, Mukaida N, Kubota Y, Kuno K, Yasumoto K, Ichimura F, Nakanishi I, Miyasaka M, Matsushima K. Human MCAF gene transfer enhances the metastatic capacity of a mouse cachectic adenocarcinoma cell line in vivo. *Pharm Res.* 12:1598-1604, 1995.
 165. Sugiyama Y, Kasahara T, Mukaida N, Matsushima K, Kitamura S. Chemokines in bronchoalveolar lavage fluid in summer-type hypersensitivity pneumonitis. *Eur Respir J.* 8:1084-1090, 1995.
 166. Teranishi Y, Mizutani H, Murata M, Shimizu M, Matsushima K. Increased spontaneous production of IL-8 in peripheral blood monocytes from the psoriatic patient: relation to focal infection and response to treatments. *J Dermatol Sci.* 10:8-15, 1995.
 167. Wakabayashi Y, Shono T, Isono M, Hori S, Matsushima K, Ono M, Kuwano M. Dual pathways of tubular morphogenesis of vascular endothelial cells by human glioma cells: vascular endothelial growth factor/basic fibroblast growth factor and interleukin-8. *Jpn J Cancer Res.* 86:1189-1197, 1995.
 168. Watanabe S, Mukaida N, Ikeda N, Akiyama M, Harada A, Nakanishi I, Nariuchi H, Watanabe Y, Matsushima K. Prevention of endotoxin shock by an antibody against leukocyte integrin beta 2 through inhibiting production and action of TNF. *Int Immunol.* 7:1037-1046, 1995.
 169. Xiu Q, Fujimura M, Nomura M, Saito M, Matsuda T, Akao N, Kondo K, Matsushima K. Bronchial hyperresponsiveness and airway neutrophil accumulation induced by interleukin-8 and the effect of the thromboxane A2 antagonist S-1452 in guinea-pigs. *Clin Exp Allergy.*

- 25:51-59, 1995.
170. Yamamoto N, Zou JP, Li XF, Takenaka H, Noda S, Fujii T, Ono S, Kobayashi Y, Mukaida N, Matsushima K, et al. Regulatory mechanisms for production of IFN-gamma and TNF by antitumor T cells or macrophages in the tumor-bearing state. *J Immunol.* 154:2281-2290, 1995.
 171. Yano S, Sone S, Nishioka Y, Mukaida N, Matsushima K, Ogura T. Differential effects of anti-inflammatory cytokines (IL-4, IL-10 and IL-13) on tumoricidal and chemotactic properties of human monocytes induced by monocyte chemotactic and activating factor. *J Leukoc Biol.* 57:303-309, 1995.
 172. Zhang Y, Harada A, Bluethmann H, Wang JB, Nakao S, Mukaida N, Matsushima K. Tumor necrosis factor (TNF) is a physiologic regulator of hematopoietic progenitor cells: increase of early hematopoietic progenitor cells in TNF receptor p55-deficient mice in vivo and potent inhibition of progenitor cell proliferation by TNF alpha in vitro. *Blood.* 86:2930-2937, 1995.
 173. Deleuran M, Buhl L, Ellingsen T, Harada A, Larsen CG, Matsushima K, Deleuran B. Localization of monocyte chemotactic and activating factor (MCAF/MCP-1) in psoriasis. *J Dermatol Sci.* 13:228-236, 1996.
 174. Fujimura M, Tsujiura M, Nomura M, Mizuguchi M, Matsuda T, Matsushima K. Sensory neuropeptides are not directly involved in bronchial hyperresponsiveness induced by interleukin-8 in guinea-pigs in vivo. *Clin Exp Allergy.* 26:357-362, 1996.
 175. Gesser B, Lund M, Lohse N, Vestergaard C, Matsushima K, Sindet-Pedersen S, Jensen SL, Thestrup-Pedersen K, Larsen CG. IL-8 induces T cell chemotaxis, suppresses IL-4, and up-regulates IL-8 production by CD4+ T cells. *J Leukoc Biol.* 59:407-411, 1996.
 176. Harada A, Mukaida N, Matsushima K. Interleukin 8 as a novel target for intervention therapy in acute inflammatory diseases. *Mol Med Today.* 2:482-489, 1996.
 177. Harada A, Mukaida N, Matsushima K. Use of Blocking Antibodies as Probes for in Vivo Functions of Chemokines. *Methods.* 10:166-174, 1996.
 178. Kurashima K, Mukaida N, Fujimura M, Schroder JM, Matsuda T, Matsushima K. Increase of chemokine levels in sputum precedes exacerbation of acute asthma attacks. *J Leukoc Biol.* 59:313-316, 1996.
 179. Mastronarde JG, He B, Monick MM, Mukaida N, Matsushima K, Hunninghake GW. Induction of interleukin (IL)-8 gene expression by respiratory syncytial virus involves activation of nuclear factor (NF)-kappa B and NF-IL-6. *J Infect Dis.* 174:262-267, 1996.
 180. Mukaida N, Ishikawa Y, Ikeda N, Fujioka N, Watanabe S, Kuno K, Matsushima K. Novel insight into molecular mechanism of endotoxin shock: biochemical analysis of LPS receptor signaling in a cell-free system targeting NF-kappaB and regulation of cytokine production/action through beta2 integrin in vivo. *J Leukoc Biol.* 59:145-151, 1996.
 181. Nagasawa T, Nakajima T, Tachibana K, Iizasa H, Bleul CC, Yoshie O, Matsushima K, Yoshida N, Springer TA, Kishimoto T. Molecular cloning and characterization of a murine pre-B-cell growth-stimulating factor/stromal cell-derived factor 1 receptor, a murine homolog of the human immunodeficiency virus 1 entry coreceptor fusin. *Proc Natl Acad Sci*

- U S A. 93:14726-14729, 1996.
182. Nakashima E, Oya A, Kubota Y, Kanada N, Matsushita R, Takeda K, Ichimura F, Kuno K, Mukaida N, Hirose K, Nakanishi I, Ujiie T, Matsushima K. A candidate for cancer gene therapy: MIP-1 alpha gene transfer to an adenocarcinoma cell line reduced tumorigenicity and induced protective immunity in immunocompetent mice. *Pharm Res.* 13:1896-1901, 1996.
 183. Ponglertnapagorn P, Oishi K, Iwagaki A, Sonoda F, Watanabe K, Nagatake T, Matsushima K, Matsumoto K. Airway interleukin-8 in elderly patients with bacterial lower respiratory tract infections. *Microbiol Immunol.* 40:177-182, 1996.
 184. Shono T, Ono M, Izumi H, Jimi SI, Matsushima K, Okamoto T, Kohno K, Kuwano M. Involvement of the transcription factor NF-kappaB in tubular morphogenesis of human microvascular endothelial cells by oxidative stress. *Mol Cell Biol.* 16:4231-4239, 1996.
 185. Su SB, Mukaida N, Matsushima K. Rapid secretion of intracellularly pre-stored interleukin-8 from rabbit platelets upon activation. *J Leukoc Biol.* 59:420-426, 1996.
 186. Su SB, Mukaida N, Wang J, Nomura H, Matsushima K. Preparation of specific polyclonal antibodies to a C-C chemokine receptor, CCR1, and determination of CCR1 expression on various types of leukocytes. *J Leukoc Biol.* 60:658-666, 1996.
 187. Usui N, Matsushima K, Pilaro AM, Longo DL, Wiltout RH. Antitumor effects of human recombinant interleukin-1 alpha and etoposide against human tumor cells: mechanism for synergism in vitro and activity in vivo. *Biotherapy.* 9:199-208, 1996.
 188. Utsunomiya I, Ito M, Watanabe K, Tsurufuji S, Matsushima K, Oh S. Infiltration of neutrophils by intrapleural injection of tumour necrosis factor, interleukin-1, and interleukin-8 in rats, and its modification by actinomycin D. *Br J Pharmacol.* 117:611-614, 1996.
 189. Wada T, Yokoyama H, Furuichi K, Kobayashi KI, Harada K, Naruto M, Su SB, Akiyama M, Mukaida N, Matsushima K. Intervention of crescentic glomerulonephritis by antibodies to monocyte chemotactic and activating factor (MCAF/MCP-1). *Faseb J.* 10:1418-1425, 1996.
 190. Wada T, Yokoyama H, Su SB, Mukaida N, Iwano M, Dohi K, Takahashi Y, Sasaki T, Furuichi K, Segawa C, Hisada Y, Ohta S, Takasawa K, Kobayashi K, Matsushima K. Monitoring urinary levels of monocyte chemotactic and activating factor reflects disease activity of lupus nephritis. *Kidney Int.* 49:761-767, 1996.
 191. Wang J, Zhang Y, Kasahara T, Harada A, Matsushima K, Mukaida N. Detection of mouse IL-8 receptor homologue expression on peripheral blood leukocytes and mature myeloid lineage cells in bone marrow. *J Leukoc Biol.* 60:372-381, 1996.
 192. Yano S, Yanagawa H, Nishioka Y, Mukaida N, Matsushima K, Sone S. T helper 2 cytokines differently regulate monocyte chemoattractant protein-1 production by human peripheral blood monocytes and alveolar macrophages. *J Immunol.* 157:2660-2665, 1996.
 193. Aihara M, Tsuchimoto D, Takizawa H, Azuma A, Wakebe H, Ohmoto Y, Imagawa K, Kikuchi M, Mukaida N, Matsushima K. Mechanisms involved in *Helicobacter pylori*-induced interleukin-8 production by a gastric cancer cell line, MKN45. *Infect Immun.* 65:3218-3224, 1997.

194. Ben-Baruch A, Grimm M, Bengali K, Evans GA, Chertov O, Wang JM, Howard OM, Mukaida N, Matsushima K, Oppenheim JJ. The differential ability of IL-8 and neutrophil-activating peptide-2 to induce attenuation of chemotaxis is mediated by their divergent capabilities to phosphorylate CXCR2 (IL-8 receptor B). *J Immunol.* 158:5927-5933, 1997.
195. Carvalho GL, Wakabayashi G, Shimazu M, Karahashi T, Yoshida M, Yamamoto S, Matsushima K, Mukaida N, Clark BD, Takabayashi T, Brandt CT, Kitajima M. Anti-interleukin-8 monoclonal antibody reduces free radical production and improves hemodynamics and survival rate in endotoxic shock in rabbits. *Surgery.* 122:60-68, 1997.
196. Fujiki F, Mukaida N, Hirose K, Ishida H, Harada A, Ohno S, Bluethmann H, Kawakami M, Akiyama M, Sone S, Matsushima K. Prevention of adenocarcinoma colon 26-induced cachexia by interleukin 10 gene transfer. *Cancer Res.* 57:94-99, 1997.
197. Fujimura M, Myou S, Nomura M, Matsuda T, Harada A, Mukaida N, Matsushima K, Nonomura A. Effect of thromboxane A2 antagonists on bronchial hyperresponsiveness induced immediately after interleukin-8 inhalation in guinea-pigs. *Br J Pharmacol.* 122:1015-1020, 1997.
198. Higashi H, Suzuki Y, Mukaida N, Takahashi N, Miyamoto D, Matsushima K. Intervention in endotoxin shock by sulfatide (I3SO3-GalCer) with a concomitant reduction in tumor necrosis factor alpha production. *Infect Immun.* 65:1223-1227, 1997.
199. Izumi S, Hirai K, Miyamasu M, Takahashi Y, Misaki Y, Takaishi T, Morita Y, Matsushima K, Ida N, Nakamura H, Kasahara T, Ito K. Expression and regulation of monocyte chemoattractant protein-1 by human eosinophils. *Eur J Immunol.* 27:816-824, 1997.
200. Jinquan T, Moller B, Storgaard M, Mukaida N, Bonde J, Grunnet N, Black FT, Larsen CG, Matsushima K, Thestrup-Pedersen K. Chemotaxis and IL-8 receptor expression in B cells from normal and HIV-infected subjects. *J Immunol.* 158:475-484, 1997.
201. Khabar KS, Al-Zoghaibi F, Al-Ahdal MN, Murayama T, Dhalla M, Mukaida N, Taha M, Al-Sedairy ST, Siddiqui Y, Kessie G, Matsushima K. The alpha chemokine, interleukin 8, inhibits the antiviral action of interferon alpha. *J Exp Med.* 186:1077-1085, 1997.
202. Khabar KS, Al-Zoghaibi F, Murayama T, Matsushima K, Mukaida N, Siddiqui Y, Dhalla M, Al-Ahdal MN. Interleukin-8 selectively enhances cytopathic effect (CPE) induced by positive-strand RNA viruses in the human WISH cell line. *Biochem Biophys Res Commun.* 235:774-778, 1997.
203. Kuno K, Iizasa H, Ohno S, Matsushima K. The exon/intron organization and chromosomal mapping of the mouse ADAMTS-1 gene encoding an ADAM family protein with TSP motifs. *Genomics.* 46:466-471, 1997.
204. Kuno K, Kanada N, Nakashima E, Fujiki F, Ichimura F, Matsushima K. Molecular cloning of a gene encoding a new type of metalloproteinase-disintegrin family protein with thrombospondin motifs as an inflammation associated gene. *J Biol Chem.* 272:556-562, 1997.
205. Kurashima K, Mukaida N, Fujimura M, Yasui M, Nakazumi Y, Matsuda T, Matsushima K. Elevated chemokine levels in bronchoalveolar lavage fluid of tuberculosis patients. *Am J*

- Respir Crit Care Med. 155:1474-1477, 1997.
206. Kurashima K, Mukaida N, Fujimura M, Yasui M, Shinagawa T, Matsuda T, Ohmoto Y, Matsushima K. A specific elevation of RANTES in bronchoalveolar lavage fluids of patients with chronic eosinophilic pneumonia. *Lab Invest.* 76:67-75, 1997.
 207. Lee LF, Haskill JS, Mukaida N, Matsushima K, Ting JP. Identification of tumor-specific paclitaxel (Taxol)-responsive regulatory elements in the interleukin-8 promoter. *Mol Cell Biol.* 17:5097-5105, 1997.
 208. Matsumori A, Furukawa Y, Hashimoto T, Yoshida A, Ono K, Shioi T, Okada M, Iwasaki A, Nishio R, Matsushima K, Sasayama S. Plasma levels of the monocyte chemotactic and activating factor/monocyte chemoattractant protein-1 are elevated in patients with acute myocardial infarction. *J Mol Cell Cardiol.* 29:419-423, 1997.
 209. Matsumoto T, Ikeda K, Mukaida N, Harada A, Matsumoto Y, Yamashita J, Matsushima K. Prevention of cerebral edema and infarct in cerebral reperfusion injury by an antibody to interleukin-8. *Lab Invest.* 77:119-125, 1997.
 210. Matsumoto T, Yokoi K, Mukaida N, Harada A, Yamashita J, Watanabe Y, Matsushima K. Pivotal role of interleukin-8 in the acute respiratory distress syndrome and cerebral reperfusion injury. *J Leukoc Biol.* 62:581-587, 1997.
 211. Miyamasu M, Masaki Y, Hirai K, Izumi S, Takaishi T, Matsushima K, Morita Y, Kasahara T. Eosinophilopoietic factors prime eosinophils for increased interleukin-8 generation. *Allergy.* 52:317-322, 1997.
 212. Murayama T, Ohara Y, Obuchi M, Khabar KS, Higashi H, Mukaida N, Matsushima K. Human cytomegalovirus induces interleukin-8 production by a human monocytic cell line, THP-1, through acting concurrently on AP-1- and NF-kappaB-binding sites of the interleukin-8 gene. *J Virol.* 71:5692-5695, 1997.
 213. Nishimura A, Akahoshi T, Takahashi M, Takagishi K, Itoman M, Kondo H, Takahashi Y, Yokoi K, Mukaida N, Matsushima K. Attenuation of monosodium urate crystal-induced arthritis in rabbits by a neutralizing antibody against interleukin-8. *J Leukoc Biol.* 62:444-449, 1997.
 214. Nishioka Y, Yano S, Fujiki F, Mukaida N, Matsushima K, Tsuruo T, Sone S. Combined therapy of multidrug-resistant human lung cancer with anti-P-glycoprotein antibody and monocyte chemoattractant protein-1 gene transduction: the possibility of immunological overcoming of multidrug resistance. *Int J Cancer.* 71:170-177, 1997.
 215. Oishi K, Sar B, Wada A, Hidaka Y, Matsumoto S, Amano H, Sonoda F, Kobayashi S, Hirayama T, Nagatake T, Matsushima K. Nitrite reductase from *Pseudomonas aeruginosa* induces inflammatory cytokines in cultured respiratory cells. *Infect Immun.* 65:2648-2655, 1997.
 216. Shioi T, Matsumori A, Kihara Y, Inoko M, Ono K, Iwanaga Y, Yamada T, Iwasaki A, Matsushima K, Sasayama S. Increased expression of interleukin-1 beta and monocyte chemotactic and activating factor/monocyte chemoattractant protein-1 in the hypertrophied and failing heart with pressure overload. *Circ Res.* 81:664-671, 1997.
 217. Sonoda Y, Kasahara T, Yamaguchi Y, Kuno K, Matsushima K, Mukaida N. Stimulation of

- interleukin-8 production by okadaic acid and vanadate in a human promyelocyte cell line, an HL-60 subline. Possible role of mitogen-activated protein kinase on the okadaic acid-induced NF-kappaB activation. *J Biol Chem.* 272:15366-15372, 1997.
218. Su S, Mukaida N, Wang J, Zhang Y, Takami A, Nakao S, Matsushima K. Inhibition of immature erythroid progenitor cell proliferation by macrophage inflammatory protein-1alpha by interacting mainly with a C-C chemokine receptor, CCR1. *Blood.* 90:605-611, 1997.
219. Sugiyama Y, Kasahara T, Mukaida N, Matsushima K, Kitamura S. Chemokines in the bronchoalveolar lavage fluid of patients with sarcoidosis. *Intern Med.* 36:856-860, 1997.
220. Tsuji H, Harada A, Mukaida N, Nakanuma Y, Bluethmann H, Kaneko S, Yamakawa K, Nakamura SI, Kobayashi KI, Matsushima K. Tumor necrosis factor receptor p55 is essential for intrahepatic granuloma formation and hepatocellular apoptosis in a murine model of bacterium-induced fulminant hepatitis. *Infect Immun.* 65:1892-1898, 1997.
221. Vestergaard C, Gesser B, Lohse N, Jensen SL, Sindet-Pedersen S, Thestrup-Pedersen K, Matsushima K, Larsen CG. Monocyte chemotactic and activating factor (MCAF/MCP-1) has an autoinductive effect in monocytes, a process regulated by IL-10. *J Dermatol Sci.* 15:14-22, 1997.
222. Wang J, Mukaida N, Zhang Y, Ito T, Nakao S, Matsushima K. Enhanced mobilization of hematopoietic progenitor cells by mouse MIP-2 and granulocyte colony-stimulating factor in mice. *J Leukoc Biol.* 62:503-509, 1997.
223. Yokoi K, Mukaida N, Harada A, Watanabe Y, Matsushima K. Prevention of endotoxemia-induced acute respiratory distress syndrome-like lung injury in rabbits by a monoclonal antibody to IL-8. *Lab Invest.* 76:375-384, 1997.
224. Zhang Y, Mukaida N, Wang J, Harada A, Akiyama M, Matsushima K. Induction of dendritic cell differentiation by granulocyte-macrophage colony-stimulating factor, stem cell factor, and tumor necrosis factor alpha in vitro from lineage phenotypes-negative c-kit+ murine hematopoietic progenitor cells. *Blood.* 90:4842-4853, 1997.
225. Aihara M, Azuma A, Takizawa H, Tsuchimoto D, Funakoshi Y, Shindo Y, Ohmoto Y, Imagawa K, Kikuchi M, Mukaida N, Matsushima K. Molecular analysis of suppression of interleukin-8 production by rebamipide in *Helicobacter pylori*-stimulated gastric cancer cell lines. *Dig Dis Sci.* 43:174S-180S, 1998.
226. Arai F, Takahashi T, Furukawa K, Matsushima K, Asakura H. Mucosal expression of interleukin-6 and interleukin-8 messenger RNA in ulcerative colitis and in Crohn's disease. *Dig Dis Sci.* 43:2071-2079, 1998.
227. Asagoe K, Yamamoto K, Takahashi A, Suzuki K, Maeda A, Nohgawa M, Harakawa N, Takano K, Mukaida N, Matsushima K, Okuma M, Sasada M. Down-regulation of CXCR2 expression on human polymorphonuclear leukocytes by TNF-alpha. *J Immunol.* 160:4518-4525, 1998.
228. Endo Y, Tabata T, Kuroda H, Tadano T, Matsushima K, Watanabe M. Induction of histidine decarboxylase in skeletal muscle in mice by electrical stimulation, prolonged walking and interleukin-1. *J Physiol.* 509 (Pt 2):587-598, 1998.
229. Fujimura M, Xiu Q, Tsujiura M, Tachibana H, Myou S, Matsuda T, Matsushima K. Role of

- leukotriene B4 in bronchial hyperresponsiveness induced by interleukin-8. *Eur Respir J*. 11:306-311, 1998.
230. Furukawa K, Takahashi T, Arai F, Matsushima K, Asakura H. Enhanced mucosal expression of interleukin-6 mRNA but not of interleukin-8 mRNA at the margin of gastric ulcer in *Helicobacter pylori*-positive gastritis. *J Gastroenterol*. 33:625-633, 1998.
231. Guo BY, Zhang SY, Mukaida N, Harada A, Kuno K, Wang JB, Sun SH, Matsushima K. CCR5 gene expression in fulminant hepatitis and DTH in mice. *World J Gastroenterol*. 4:14-18, 1998.
232. Higashi H, Mukaida N, Harada A, Watanabe S, Ikeda N, Suzuki Y, Matsushima K. Prevention of endotoxin shock through targeting leukocyte adhesion molecules. *Prog Clin Biol Res*. 397:327-334, 1998.
233. Hu H, Shioda T, Hori T, Moriya C, Kato A, Sakai Y, Matsushima K, Uchiyama T, Nagai Y. Dissociation of ligand-induced internalization of CXCR-4 from its co-receptor activity for HIV-1 Env-mediated membrane fusion. *Arch Virol*. 143:851-861, 1998.
234. Kasahara T, Oda T, Hatake K, Akiyama M, Mukaida N, Matsushima K. Interleukin-8 and monocyte chemotactic protein-1 production by a human glioblastoma cell line, T98G in coculture with monocytes: involvement of monocyte-derived interleukin-1alpha. *Eur Cytokine Netw*. 9:47-55, 1998.
235. Kasahara Y, Kimura H, Kurosu K, Sugito K, Mukaida N, Matsushima K, Kuriyama T. MCAF/MCP-1 protein expression in a rat model for pulmonary hypertension induced by monocrotaline. *Chest*. 114:67S, 1998.
236. Kimura H, Kasahara Y, Kurosu K, Sugito K, Takiguchi Y, Terai M, Mikata A, Natsume M, Mukaida N, Matsushima K, Kuriyama T. Alleviation of monocrotaline-induced pulmonary hypertension by antibodies to monocyte chemotactic and activating factor/monocyte chemoattractant protein-1. *Lab Invest*. 78:571-581, 1998.
237. Kuno K, Matsushima K. ADAMTS-1 protein anchors at the extracellular matrix through the thrombospondin type I motifs and its spacing region. *J Biol Chem*. 273:13912-13917, 1998.
238. Mastronarde JG, Monick MM, Mukaida N, Matsushima K, Hunninghake GW. Activator protein-1 is the preferred transcription factor for cooperative interaction with nuclear factor-kappaB in respiratory syncytial virus-induced interleukin-8 gene expression in airway epithelium. *J Infect Dis*. 177:1275-1281, 1998.
239. Miyamasu M, Misaki Y, Izumi S, Takaishi T, Morita Y, Nakamura H, Matsushima K, Kasahara T, Hirai K. Glucocorticoids inhibit chemokine generation by human eosinophils. *J Allergy Clin Immunol*. 101:75-83, 1998.
240. Mori N, Mukaida N, Ballard DW, Matsushima K, Yamamoto N. Human T-cell leukemia virus type I Tax transactivates human interleukin 8 gene through acting concurrently on AP-1 and nuclear factor-kappaB-like sites. *Cancer Res*. 58:3993-4000, 1998.
241. Mukaida N, Harada A, Matsushima K. Interleukin-8 (IL-8) and monocyte chemotactic and activating factor (MCAF/MCP-1), chemokines essentially involved in inflammatory and immune reactions. *Cytokine Growth Factor Rev*. 9:9-23, 1998.
242. Mukaida N, Matsumoto T, Yokoi K, Harada A, Matsushima K. Inhibition of

- neutrophil-mediated acute inflammation injury by an antibody against interleukin-8 (IL-8). *Inflamm Res.* 47 Suppl 3:S151-157, 1998.
243. Murayama T, Mukaida N, Khabar KS, Matsushima K. Potential involvement of IL-8 in the pathogenesis of human cytomegalovirus infection. *J Leukoc Biol.* 64:62-67, 1998.
244. Nakashima E, Kubota Y, Matsushita R, Ozaki E, Ichimura F, Kawahara S, Nakanishi I, Kuno K, Matsushima K. Synergistic antitumor interaction of human monocyte chemoattractant protein-1 gene transfer and modulator for tumor-infiltrating macrophages. *Pharm Res.* 15:685-689, 1998.
245. Okada M, Matsumori A, Ono K, Furukawa Y, Shioi T, Iwasaki A, Matsushima K, Sasayama S. Cyclic stretch upregulates production of interleukin-8 and monocyte chemoattractant protein-1 in human endothelial cells. *Arterioscler Thromb Vasc Biol.* 18:894-901, 1998.
246. Osman MO, Kristensen JU, Jacobsen NO, Lausten SB, Deleuran B, Deleuran M, Gesser B, Matsushima K, Larsen CG, Jensen SL. A monoclonal anti-interleukin 8 antibody (WS-4) inhibits cytokine response and acute lung injury in experimental severe acute necrotising pancreatitis in rabbits. *Gut.* 43:232-239, 1998.
247. Roger T, Out T, Mukaida N, Matsushima K, Jansen H, Lutter R. Enhanced AP-1 and NF-kappaB activities and stability of interleukin 8 (IL-8) transcripts are implicated in IL-8 mRNA superinduction in lung epithelial H292 cells. *Biochem J.* 330 (Pt 1):429-435, 1998.
248. Shimoya K, Matsuzaki N, Sawai K, Saji F, Murata Y, Yasumoto K, Su S, Mukaida N, Matsushima K. Regulation of placental monocyte chemoattractant and activating factor during pregnancy and chorioamnionitis. *Mol Hum Reprod.* 4:393-400, 1998.
249. Sonoda Y, Mukaida N, Wang JB, Shimada-Hiratsuka M, Naito M, Kasahara T, Harada A, Inoue M, Matsushima K. Physiologic regulation of postovulatory neutrophil migration into vagina in mice by a C-X-C chemokine(s). *J Immunol.* 160:6159-6165, 1998.
250. Tachibana K, Hirota S, Iizasa H, Yoshida H, Kawabata K, Kataoka Y, Kitamura Y, Matsushima K, Yoshida N, Nishikawa S, Kishimoto T, Nagasawa T. The chemokine receptor CXCR4 is essential for vascularization of the gastrointestinal tract. *Nature.* 393:591-594, 1998.
251. Tozawa Y, Ueki A, Manabe S, Matsushima K. Stress-induced increase in urinary isatin excretion in rats: reversal by both dexamethasone and alpha-methyl-P-tyrosine. *Biochem Pharmacol.* 56:1041-1046, 1998.
252. Wang J, Harada A, Matsushita S, Matsumi S, Zhang Y, Shioda T, Nagai Y, Matsushima K. IL-4 and a glucocorticoid up-regulate CXCR4 expression on human CD4+ T lymphocytes and enhance HIV-1 replication. *J Leukoc Biol.* 64:642-649, 1998.
253. Yamada T, Hisanaga M, Nakajima Y, Kanehiro H, Watanabe A, Ohyama T, Nishio K, Sho M, Nagao M, Harada A, Matsushima K, Nakano H. Serum interleukin-6, interleukin-8, hepatocyte growth factor, and nitric oxide changes during thoracic surgery. *World J Surg.* 22:783-790, 1998.
254. Yokoyama H, Wada T, Furuichi K, Segawa C, Shimizu M, Kobayashi K, Su S, Mukaida N, Matsushima K. Urinary levels of chemokines (MCAF/MCP-1, IL-8) reflect distinct disease

- activities and phases of human IgA nephropathy. *J Leukoc Biol.* 63:493-499, 1998.
255. Yoneyama H, Harada A, Imai T, Baba M, Yoshie O, Zhang Y, Higashi H, Murai M, Asakura H, Matsushima K. Pivotal role of TARC, a CC chemokine, in bacteria-induced fulminant hepatic failure in mice. *J Clin Invest.* 102:1933-1941, 1998.
256. Yoshimine H, Oishi K, Nagatake T, Matsushima K, Omura T. Inhibition of neutrophil-derived IL-8 production caused by EM (erythromycin) 201 derivative: modification of the activities by the 5-position desosamine side chain. *Jpn J Antibiot.* 51 Suppl:134-136, 1998.
257. Zhang Y, Harada A, Wang JB, Zhang YY, Hashimoto S, Naito M, Matsushima K. Bifurcated dendritic cell differentiation in vitro from murine lineage phenotype-negative c-kit⁺ bone marrow hematopoietic progenitor cells. *Blood.* 92:118-128, 1998.
258. Asahi K, Mizutani H, Tanaka M, Miura M, Yamanaka K, Matsushima K, Nakashima K, Shimizu M. Intradermal transfer of caspase-1 (CASP1) DNA into mouse dissects: role of CASP1 in interleukin-1beta associated skin inflammation and apoptotic cell death. *J Dermatol Sci.* 21:49-58, 1999.
259. Betsuyaku T, Liu F, Senior RM, Haug JS, Brown EJ, Jones SL, Matsushima K, Link DC. A functional granulocyte colony-stimulating factor receptor is required for normal chemoattractant-induced neutrophil activation. *J Clin Invest.* 103:825-832, 1999.
260. De Benedetti F, Pignatti P, Bernasconi S, Gerloni V, Matsushima K, Caporali R, Montecucco CM, Sozzani S, Fantini F, Martini A. Interleukin 8 and monocyte chemoattractant protein-1 in patients with juvenile rheumatoid arthritis. Relation to onset types, disease activity, and synovial fluid leukocytes. *J Rheumatol.* 26:425-431, 1999.
261. Fujimura M, Myou S, Nomura M, Mizuguchi M, Matsuda T, Harada A, Mukaida N, Matsushima K, Nonomura A. Interleukin-8 inhalation directly provokes bronchoconstriction in guinea pigs. *Allergy.* 54:386-391, 1999.
262. Furukawa Y, Matsumori A, Ohashi N, Shioi T, Ono K, Harada A, Matsushima K, Sasayama S. Anti-monocyte chemoattractant protein-1/monocyte chemoattractant and activating factor antibody inhibits neointimal hyperplasia in injured rat carotid arteries. *Circ Res.* 84:306-314, 1999.
263. Hashimoto S, Suzuki T, Dong HY, Nagai S, Yamazaki N, Matsushima K. Serial analysis of gene expression in human monocyte-derived dendritic cells. *Blood.* 94:845-852, 1999.
264. Hashimoto S, Suzuki T, Dong HY, Yamazaki N, Matsushima K. Serial analysis of gene expression in human monocytes and macrophages. *Blood.* 94:837-844, 1999.
265. Imai T, Nagira M, Takagi S, Kakizaki M, Nishimura M, Wang J, Gray PW, Matsushima K, Yoshie O. Selective recruitment of CCR4-bearing Th2 cells toward antigen-presenting cells by the CC chemokines thymus and activation-regulated chemokine and macrophage-derived chemokine. *Int Immunol.* 11:81-88, 1999.
266. Inadera H, Egashira K, Takemoto M, Ouchi Y, Matsushima K. Increase in circulating levels of monocyte chemoattractant protein-1 with aging. *J Interferon Cytokine Res.* 19:1179-1182, 1999.
267. Karima R, Matsumoto S, Higashi H, Matsushima K. The molecular pathogenesis of

- endotoxic shock and organ failure. *Mol Med Today*. 5:123-132, 1999.
268. Kuno K, Terashima Y, Matsushima K. ADAMTS-1 is an active metalloproteinase associated with the extracellular matrix. *J Biol Chem*. 274:18821-18826, 1999.
269. Miyamasu M, Yamaguchi M, Nakajima T, Misaki Y, Morita Y, Matsushima K, Yamamoto K, Hirai K. Th1-derived cytokine IFN-gamma is a potent inhibitor of eotaxin synthesis in vitro. *Int Immunol*. 11:1001-1004, 1999.
270. Mori N, Oishi K, Sar B, Mukaida N, Nagatake T, Matsushima K, Yamamoto N. Essential role of transcription factor nuclear factor-kappaB in regulation of interleukin-8 gene expression by nitrite reductase from *Pseudomonas aeruginosa* in respiratory epithelial cells. *Infect Immun*. 67:3872-3878, 1999.
271. Murai M, Yoneyama H, Harada A, Yi Z, Vestergaard C, Guo B, Suzuki K, Asakura H, Matsushima K. Active participation of CCR5(+)/CD8(+) T lymphocytes in the pathogenesis of liver injury in graft-versus-host disease. *J Clin Invest*. 104:49-57, 1999.
272. Natsume M, Tsuji H, Harada A, Akiyama M, Yano T, Ishikura H, Nakanishi I, Matsushima K, Kaneko S, Mukaida N. Attenuated liver fibrosis and depressed serum albumin levels in carbon tetrachloride-treated IL-6-deficient mice. *J Leukoc Biol*. 66:601-608, 1999.
273. Nokihara H, Nishioka Y, Yano S, Mukaida N, Matsushima K, Tsuruo T, Sone S. Monocyte chemoattractant protein-1 gene modification of multidrug-resistant human lung cancer enhances antimetastatic effect of therapy with anti-P-glycoprotein antibody in SCID mice. *Int J Cancer*. 80:773-780, 1999.
274. Ogata M, Zhang Y, Wang Y, Itakura M, Zhang YY, Harada A, Hashimoto S, Matsushima K. Chemotactic response toward chemokines and its regulation by transforming growth factor-beta1 of murine bone marrow hematopoietic progenitor cell-derived different subset of dendritic cells. *Blood*. 93:3225-3232, 1999.
275. Ono K, Matsumori A, Furukawa Y, Igata H, Shioi T, Matsushima K, Sasayama S. Prevention of myocardial reperfusion injury in rats by an antibody against monocyte chemoattractant and activating factor/monocyte chemoattractant protein-1. *Lab Invest*. 79:195-203, 1999.
276. Sansonetti PJ, Arondel J, Huerre M, Harada A, Matsushima K. Interleukin-8 controls bacterial transepithelial translocation at the cost of epithelial destruction in experimental shigellosis. *Infect Immun*. 67:1471-1480, 1999.
277. Sar B, Oishi K, Matsushima K, Nagatake T. Induction of interleukin 8 (IL-8) production by *Pseudomonas* nitrite reductase in human alveolar macrophages and epithelial cells. *Microbiol Immunol*. 43:409-417, 1999.
278. Sar B, Oishi K, Wada A, Hirayama T, Matsushima K, Nagatake T. Nitrite reductase from *Pseudomonas aeruginosa* released by antimicrobial agents and complement induces interleukin-8 production in bronchial epithelial cells. *Antimicrob Agents Chemother*. 43:794-801, 1999.
279. Shi Q, Le X, Abbruzzese JL, Wang B, Mujaida N, Matsushima K, Huang S, Xiong Q, Xie K. Cooperation between transcription factor AP-1 and NF-kappaB in the induction of interleukin-8 in human pancreatic adenocarcinoma cells by hypoxia. *J Interferon Cytokine Res*. 19:1363-1371, 1999.

280. Suzuki N, Nakajima A, Yoshino S, Matsushima K, Yagita H, Okumura K. Selective accumulation of CCR5+ T lymphocytes into inflamed joints of rheumatoid arthritis. *Int Immunol.* 11:553-559, 1999.
281. Tadano T, Namioka M, Nakagawasai O, Tan-No K, Matsushima K, Endo Y, Kisara K. Induction of nociceptive responses by intrathecal injection of interleukin-1 in mice. *Life Sci.* 65:255-261, 1999.
282. Takahashi Y, Kasahara T, Sawai T, Rikimaru A, Mukaida N, Matsushima K, Sasaki T. The participation of IL-8 in the synovial lesions at an early stage of rheumatoid arthritis. *Tohoku J Exp Med.* 188:75-87, 1999.
283. Takizawa H, Ohtoshi T, Kawasaki S, Kohyama T, Desaki M, Kasama T, Kobayashi K, Nakahara K, Yamamoto K, Matsushima K, Kudoh S. Diesel exhaust particles induce NF-kappa B activation in human bronchial epithelial cells in vitro: importance in cytokine transcription. *J Immunol.* 162:4705-4711, 1999.
284. Terai M, Jibiki T, Harada A, Terashima Y, Yasukawa K, Tateno S, Hamada H, Oana S, Niimi H, Matsushima K. Dramatic decrease of circulating levels of monocyte chemoattractant protein-1 in Kawasaki disease after gamma globulin treatment. *J Leukoc Biol.* 65:566-572, 1999.
285. Tsuji H, Mukaida N, Harada A, Kaneko S, Matsushita E, Nakanuma Y, Tsutsui H, Okamura H, Nakanishi K, Tagawa Y, Iwakura Y, Kobayashi K, Matsushima K. Alleviation of lipopolysaccharide-induced acute liver injury in Propionibacterium acnes-primed IFN-gamma-deficient mice by a concomitant reduction of TNF-alpha, IL-12, and IL-18 production. *J Immunol.* 162:1049-1055, 1999.
286. Verma MJ, Mukaida N, Vollmer-Conna U, Matsushima K, Lloyd A, Wakefield D. Endotoxin-induced uveitis is partially inhibited by anti-IL-8 antibody treatment. *Invest Ophthalmol Vis Sci.* 40:2465-2470, 1999.
287. Vestergaard C, Yoneyama H, Murai M, Nakamura K, Tamaki K, Terashima Y, Imai T, Yoshie O, Irimura T, Mizutani H, Matsushima K. Overproduction of Th2-specific chemokines in NC/Nga mice exhibiting atopic dermatitis-like lesions. *J Clin Invest.* 104:1097-1105, 1999.
288. Wada T, Furuichi K, Sakai N, Shimizu M, Segawa C, Kobayashi K, Mukaida N, Kasahara T, Matsushima K, Yokoyama H. Eotaxin contributes to renal interstitial eosinophilia. *Nephrol Dial Transplant.* 14:76-80, 1999.
289. Wada T, Furuichi K, Segawa-Takaeda C, Shimizu M, Sakai N, Takeda SI, Takasawa K, Kida H, Kobayashi KI, Mukaida N, Ohmoto Y, Matsushima K, Yokoyama H. MIP-1alpha and MCP-1 contribute to crescents and interstitial lesions in human crescentic glomerulonephritis. *Kidney Int.* 56:995-1003, 1999.
290. Xu L, Xie K, Mukaida N, Matsushima K, Fidler IJ. Hypoxia-induced elevation in interleukin-8 expression by human ovarian carcinoma cells. *Cancer Res.* 59:5822-5829, 1999.
291. Yamada T, Hisanaga M, Nakajima Y, Kanehiro H, Aomatsu Y, Ko S, Kin T, Nishio K, Sho M, Nagao M, Harada A, Matsushima K, Nakano H. The serum interleukin 8 level reflects hepatic mitochondrial redox state in hyperthermochemohypoxic isolated liver perfusion with

- use of a venovenous bypass. *Surgery*. 125:304-314, 1999.
292. Yamada Y, Karasaki H, Matsushima K, Lee GH, Ogawa K. Expression of an IL-1 receptor antagonist during mouse hepatocarcinogenesis demonstrated by differential display analysis. *Lab Invest*. 79:1059-1067, 1999.
293. Zhang Y, Zhang YY, Ogata M, Chen P, Harada A, Hashimoto S, Matsushima K. Transforming growth factor-beta1 polarizes murine hematopoietic progenitor cells to generate Langerhans cell-like dendritic cells through a monocyte/macrophage differentiation pathway. *Blood*. 93:1208-1220, 1999.
294. Abe S, Nakamura H, Inoue S, Takeda H, Saito H, Kato S, Mukaida N, Matsushima K, Tomoike H. Interleukin-8 gene repression by clarithromycin is mediated by the activator protein-1 binding site in human bronchial epithelial cells. *Am J Respir Cell Mol Biol*. 22:51-60, 2000.
295. Amano H, Yamamoto H, Senba M, Oishi K, Suzuki S, Fukushima K, Mukaida N, Matsushima K, Eguchi K, Nagatake T. Impairment of endotoxin-induced macrophage inflammatory protein 2 gene expression in alveolar macrophages in streptozotocin-induced diabetes in mice. *Infect Immun*. 68:2925-2929, 2000.
296. Ato M, Iwabuchi K, Matsuki N, Mukaida N, Iwabuchi C, Takahashi A, Takayanagi T, Dondog EA, Hatakeyama S, Ishikura H, Kato M, Negishi I, Nishihori H, Watano K, Ogasawara K, Matsushima K, Onoe K. Delayed clearance of zymosan-induced granuloma and depressed phagocytosis of macrophages with concomitant up-regulated kinase activities of Src-family in a human monocyte chemoattractant protein-1 transgenic mouse. *Immunobiology*. 201:432-449, 2000.
297. Ellingsen T, Elling P, Olson A, Elling H, Baandrup U, Matsushima K, Deleuran B, Stengaard-Pedersen K. Monocyte chemoattractant protein 1 (MCP-1) in temporal arteritis and polymyalgia rheumatica. *Ann Rheum Dis*. 59:775-780, 2000.
298. Furuichi K, Wada T, Sakai N, Iwata Y, Yoshimoto K, Shimizu M, Kobayashi K, Takasawa K, Kida H, Takeda SI, Mukaida N, Matsushima K, Yokoyama H. Distinct expression of CCR1 and CCR5 in glomerular and interstitial lesions of human glomerular diseases. *Am J Nephrol*. 20:291-299, 2000.
299. Hashimoto SI, Suzuki T, Nagai S, Yamashita T, Toyoda N, Matsushima K. Identification of genes specifically expressed in human activated and mature dendritic cells through serial analysis of gene expression. *Blood*. 96:2206-2214, 2000.
300. Hirao M, Onai N, Hiroishi K, Watkins SC, Matsushima K, Robbins PD, Lotze MT, Tahara H. CC chemokine receptor-7 on dendritic cells is induced after interaction with apoptotic tumor cells: critical role in migration from the tumor site to draining lymph nodes. *Cancer Res*. 60:2209-2217, 2000.
301. Horiguchi H, Harada A, Oguma E, Sato M, Homma Y, Kayama F, Fukushima M, Matsushima K. Cadmium-induced acute hepatic injury is exacerbated in human interleukin-8 transgenic mice. *Toxicol Appl Pharmacol*. 163:231-239, 2000.
302. Iguchi A, Kitajima I, Yamakuchi M, Ueno S, Aikou T, Kubo T, Matsushima K, Mukaida N, Maruyama I. PEA3 and AP-1 are required for constitutive IL-8 gene expression in hepatoma

- cells. *Biochem Biophys Res Commun.* 279:166-171, 2000.
303. Iguchi H, Ono M, Matsushima K, Kuwano M. Overproduction of IL-8 results in suppression of bone metastasis by lung cancer cells in vivo. *Int J Oncol.* 17:329-333, 2000.
304. Inadera H, Hashimoto S, Dong HY, Suzuki T, Nagai S, Yamashita T, Toyoda N, Matsushima K. WISP-2 as a novel estrogen-responsive gene in human breast cancer cells. *Biochem Biophys Res Commun.* 275:108-114, 2000.
305. Inadera H, Sekiya T, Yoshimura T, Matsushima K. Molecular analysis of the inhibition of monocyte chemoattractant protein-1 gene expression by estrogens and xenoestrogens in MCF-7 cells. *Endocrinology.* 141:50-59, 2000.
306. Ishii M, Hashimoto S, Tsutsumi S, Wada Y, Matsushima K, Kodama T, Aburatani H. Direct comparison of GeneChip and SAGE on the quantitative accuracy in transcript profiling analysis. *Genomics.* 68:136-143, 2000.
307. Kakio T, Matsumori A, Ono K, Ito H, Matsushima K, Sasayama S. Roles and relationship of macrophages and monocyte chemoattractant and activating factor/monocyte chemoattractant protein-1 in the ischemic and reperfused rat heart. *Lab Invest.* 80:1127-1136, 2000.
308. Kato N, Yoshida H, Kioko Ono-Nita S, Kato J, Goto T, Otsuka M, Lan K, Matsushima K, Shiratori Y, Omata M. Activation of intracellular signaling by hepatitis B and C viruses: C-viral core is the most potent signal inducer. *Hepatology.* 32:405-412, 2000.
309. Kuno K, Okada Y, Kawashima H, Nakamura H, Miyasaka M, Ohno H, Matsushima K. ADAMTS-1 cleaves a cartilage proteoglycan, aggrecan. *FEBS Lett.* 478:241-245, 2000.
310. Lee LF, Hellendall RP, Wang Y, Haskill JS, Mukaida N, Matsushima K, Ting JP. IL-8 reduced tumorigenicity of human ovarian cancer in vivo due to neutrophil infiltration. *J Immunol.* 164:2769-2775, 2000.
311. Matsugo S, Kitagawa T, Minami S, Esashi Y, Oomura Y, Tokumaru S, Kojo S, Matsushima K, Sasaki K. Age-dependent changes in lipid peroxide levels in peripheral organs, but not in brain, in senescence-accelerated mice. *Neurosci Lett.* 278:105-108, 2000.
312. Matsushima K. Chemokines. Introduction. *Springer Semin Immunopathol.* 22:321-328, 2000.
313. Miyamasu M, Misiaki Y, Yamaguchi M, Yamamoto K, Morita Y, Matsushima K, Nakajima T, Hirai K. Regulation of human eotaxin generation by Th1-/Th2-derived cytokines. *Int Arch Allergy Immunol.* 122 Suppl 1:54-58, 2000.
314. Murayama T, Mukaida N, Sadanari H, Yamaguchi N, Khabar KS, Tanaka J, Matsushima K, Mori S, Eizuru Y. The immediate early gene 1 product of human cytomegalovirus is sufficient for up-regulation of interleukin-8 gene expression. *Biochem Biophys Res Commun.* 279:298-304, 2000.
315. Murphy PM, Baggiolini M, Charo IF, Hebert CA, Horuk R, Matsushima K, Miller LH, Oppenheim JJ, Power CA. International union of pharmacology. XXII. Nomenclature for chemokine receptors. *Pharmacol Rev.* 52:145-176, 2000.
316. Narumi S, Yoneyama H, Inadera H, Nishioji K, Itoh Y, Okanoue T, Matsushima K. TNF-alpha is a potent inducer for IFN-inducible protein-10 in hepatocytes and unaffected by GM-CSF in vivo, in contrast to IL-1beta and IFN-gamma. *Cytokine.* 12:1007-1016, 2000.

317. Nishihori H, Tsuji H, Wang H, Tahara H, Akiyama M, Ogawa Y, Matsushima K, Iwakura Y, Mukaida N. Participation of endogenously produced interferon gamma in interleukin 4-mediated tumor rejection. *Hum Gene Ther.* 11:659-668, 2000.
318. Nokihara H, Yanagawa H, Nishioka Y, Yano S, Mukaida N, Matsushima K, Sone S. Natural killer cell-dependent suppression of systemic spread of human lung adenocarcinoma cells by monocyte chemoattractant protein-1 gene transfection in severe combined immunodeficient mice. *Cancer Res.* 60:7002-7007, 2000.
319. Oishi K, Hayano M, Yoshimine H, Tugume SB, Kebba A, Mugerwa R, Mugenyi P, Kumatori A, Matsushima K, Nagatake T. Expression of chemokine receptors on CD4+ T cells in peripheral blood from HIV-infected individuals in Uganda. *J Interferon Cytokine Res.* 20:597-602, 2000.
320. Onai N, Zhang Y, Yoneyama H, Kitamura T, Ishikawa S, Matsushima K. Impairment of lymphopoiesis and myelopoiesis in mice reconstituted with bone marrow-hematopoietic progenitor cells expressing SDF-1-intrakine. *Blood.* 96:2074-2080, 2000.
321. Osawa Y, Hachiya M, Araki S, Kusama T, Matsushima K, Aoki Y, Akashi M. IL-1 induces expression of p21(WAF1) independently of p53 in high-passage human embryonic fibroblasts WI38. *J Biochem (Tokyo).* 127:883-893, 2000.
322. Ostergaard C, Yieng-Kow RV, Larsen CG, Mukaida N, Matsushima K, Benfield T, Frimodt-Moller N, Espersen F, Kharazmi A, Lundgren JD. Treatment with a monoclonal antibody to IL-8 attenuates the pleocytosis in experimental pneumococcal meningitis in rabbits when given intravenously, but not intracisternally. *Clin Exp Immunol.* 122:207-211, 2000.
323. Sar B, Oishi K, Wada A, Hirayama T, Matsushima K, Nagatake T. Induction of monocyte chemoattractant protein-1 (MCP-1) production by *Pseudomonas* nitrite reductase in human pulmonary type II epithelial-like cells. *Microb Pathog.* 28:17-23, 2000.
324. Sekiya T, Miyamasu M, Imanishi M, Yamada H, Nakajima T, Yamaguchi M, Fujisawa T, Pawankar R, Sano Y, Ohta K, Ishii A, Morita Y, Yamamoto K, Matsushima K, Yoshie O, Hirai K. Inducible expression of a Th2-type CC chemokine thymus- and activation-regulated chemokine by human bronchial epithelial cells. *J Immunol.* 165:2205-2213, 2000.
325. Shindo T, Kurihara H, Kuno K, Yokoyama H, Wada T, Kurihara Y, Imai T, Wang Y, Ogata M, Nishimatsu H, Moriyama N, Oh-hashii Y, Morita H, Ishikawa T, Nagai R, Yazaki Y, Matsushima K. ADAMTS-1: a metalloproteinase-disintegrin essential for normal growth, fertility, and organ morphology and function. *J Clin Invest.* 105:1345-1352, 2000.
326. Suzuki T, Hashimoto S, Toyoda N, Nagai S, Yamazaki N, Dong HY, Sakai J, Yamashita T, Nukiwa T, Matsushima K. Comprehensive gene expression profile of LPS-stimulated human monocytes by SAGE. *Blood.* 96:2584-2591, 2000.
327. Takizawa H, Ohtoshi T, Kawasaki S, Abe S, Sugawara I, Nakahara K, Matsushima K, Kudoh S. Diesel exhaust particles activate human bronchial epithelial cells to express inflammatory mediators in the airways: a review. *Respirology.* 5:197-203, 2000.
328. Tamaru M, Nishioji K, Kobayashi Y, Watanabe Y, Itoh Y, Okanoue T, Murai M, Matsushima K, Narumi S. Liver-infiltrating T lymphocytes are attracted selectively by IFN-inducible

- protein-10. *Cytokine*. 12:299-308, 2000.
329. Tokuda A, Itakura M, Onai N, Kimura H, Kuriyama T, Matsushima K. Pivotal role of CCR1-positive leukocytes in bleomycin-induced lung fibrosis in mice. *J Immunol*. 164:2745-2751, 2000.
330. Ueno T, Toi M, Saji H, Muta M, Bando H, Kuroi K, Koike M, Inadera H, Matsushima K. Significance of macrophage chemoattractant protein-1 in macrophage recruitment, angiogenesis, and survival in human breast cancer. *Clin Cancer Res*. 6:3282-3289, 2000.
331. Usui M, Egashira K, Tomita H, Koyanagi M, Katoh M, Shimokawa H, Takeya M, Yoshimura T, Matsushima K, Takeshita A. Important role of local angiotensin II activity mediated via type 1 receptor in the pathogenesis of cardiovascular inflammatory changes induced by chronic blockade of nitric oxide synthesis in rats. *Circulation*. 101:305-310, 2000.
332. Vestergaard C, Bang K, Gesser B, Yoneyama H, Matsushima K, Larsen CG. A Th2 chemokine, TARC, produced by keratinocytes may recruit CLA+CCR4+ lymphocytes into lesional atopic dermatitis skin. *J Invest Dermatol*. 115:640-646, 2000.
333. Vestergaard C, Yoneyama H, Matsushima K. The NC/Nga mouse: a model for atopic dermatitis. *Mol Med Today*. 6:209-210, 2000.
334. Wada T, Furuichi K, Sakai N, Iwata Y, Yoshimoto K, Shimizu M, Kobayashi K, Mukaida N, Matsushima K, Yokoyama H. A new anti-inflammatory compound, FR167653, ameliorates crescentic glomerulonephritis in Wistar-Kyoto rats. *J Am Soc Nephrol*. 11:1534-1541, 2000.
335. Wada T, Furuichi K, Sakai N, Iwata Y, Yoshimoto K, Shimizu M, Takeda SI, Takasawa K, Yoshimura M, Kida H, Kobayashi KI, Mukaida N, Naito T, Matsushima K, Yokoyama H. Up-regulation of monocyte chemoattractant protein-1 in tubulointerstitial lesions of human diabetic nephropathy. *Kidney Int*. 58:1492-1499, 2000.
336. Yamamoto J, Adachi Y, Onoue Y, Adachi YS, Okabe Y, Itazawa T, Toyoda M, Seki T, Morohashi M, Matsushima K, Miyawaki T. Differential expression of the chemokine receptors by the Th1- and Th2-type effector populations within circulating CD4+ T cells. *J Leukoc Biol*. 68:568-574, 2000.
337. Yamashita T, Hashimoto S, Kaneko S, Nagai S, Toyoda N, Suzuki T, Kobayashi K, Matsushima K. Comprehensive gene expression profile of a normal human liver. *Biochem Biophys Res Commun*. 269:110-116, 2000.
338. Yokoyama A, Kohno N, Ito M, Abe M, Hiwada K, Yamada H, Matsushima K, Hirai K. Eotaxin levels in pleural effusions: comparison with monocyte chemoattractant protein-1 and IL-8. *Intern Med*. 39:547-552, 2000.
339. Yoneyama H, Kawasaki S, Matsushima K. Regulation of Th1 and Th2 immune responses by chemokines. *Springer Semin Immunopathol*. 22:329-344, 2000.
340. Zhang Y, Wang Y, Ogata M, Hashimoto S, Onai N, Matsushima K. Development of dendritic cells in vitro from murine fetal liver-derived lineage phenotype-negative c-kit(+) hematopoietic progenitor cells. *Blood*. 95:138-146, 2000.
341. de Wynter EA, Heyworth CM, Mukaida N, Matsushima K, Testa NG. NOD/SCID repopulating cells but not LTC-IC are enriched in human CD34+ cells expressing the CCR1 chemokine receptor. *Leukemia*. 15:1092-1101, 2001.

342. Furuichi K, Wada T, Iwata Y, Sakai N, Yoshimoto K, Shimizu M, Kobayashi K, Takasawa K, Kida H, Takeda S, Matsushima K, Yokoyama H. Upregulation of fractalkine in human crescentic glomerulonephritis. *Nephron*. 87:314-320, 2001.
343. Ghia P, Transidico P, Veiga JP, Schaniel C, Sallusto F, Matsushima K, Sallan SE, Rolink AG, Mantovani A, Nadler LM, Cardoso AA. Chemoattractants MDC and TARC are secreted by malignant B-cell precursors following CD40 ligation and support the migration of leukemia-specific T cells. *Blood*. 98:533-540, 2001.
344. Hase K, Tani K, Shimizu T, Ohmoto Y, Matsushima K, Sone S. Increased CCR4 expression in active systemic lupus erythematosus. *J Leukoc Biol*. 70:749-755, 2001.
345. Iikura M, Miyamasu M, Yamaguchi M, Kawasaki H, Matsushima K, Kitaura M, Morita Y, Yoshie O, Yamamoto K, Hirai K. Chemokine receptors in human basophils: inducible expression of functional CXCR4. *J Leukoc Biol*. 70:113-120, 2001.
346. Ishikawa S, Sato T, Abe M, Nagai S, Onai N, Yoneyama H, Zhang Y, Suzuki T, Hashimoto S, Shirai T, Lipp M, Matsushima K. Aberrant high expression of B lymphocyte chemokine (BLC/CXCL13) by C11b+CD11c+ dendritic cells in murine lupus and preferential chemotaxis of B1 cells towards BLC. *J Exp Med*. 193:1393-1402, 2001.
347. Itakura M, Tokuda A, Kimura H, Nagai S, Yoneyama H, Onai N, Ishikawa S, Kuriyama T, Matsushima K. Blockade of secondary lymphoid tissue chemokine exacerbates *Propionibacterium acnes*-induced acute lung inflammation. *J Immunol*. 166:2071-2079, 2001.
348. Kakinuma T, Nakamura K, Wakugawa M, Mitsui H, Tada Y, Saeki H, Torii H, Asahina A, Onai N, Matsushima K, Tamaki K. Thymus and activation-regulated chemokine in atopic dermatitis: Serum thymus and activation-regulated chemokine level is closely related with disease activity. *J Allergy Clin Immunol*. 107:535-541, 2001.
349. Kasahara T, Matsushima K. Macrophage signaling, apoptosis, lectins and leukocyte trafficking. *Trends Immunol*. 22:593-594, 2001.
350. Katou F, Ohtani H, Nakayama T, Ono K, Matsushima K, Saaristo A, Nagura H, Yoshie O, Motegi K. Macrophage-derived chemokine (MDC/CCL22) and CCR4 are involved in the formation of T lymphocyte-dendritic cell clusters in human inflamed skin and secondary lymphoid tissue. *Am J Pathol*. 158:1263-1270, 2001.
351. Kawasaki S, Takizawa H, Yoneyama H, Nakayama T, Fujisawa R, Izumizaki M, Imai T, Yoshie O, Homma I, Yamamoto K, Matsushima K. Intervention of thymus and activation-regulated chemokine attenuates the development of allergic airway inflammation and hyperresponsiveness in mice. *J Immunol*. 166:2055-2062, 2001.
352. Kimura H, Okada O, Tanabe N, Tanaka Y, Terai M, Takiguchi Y, Masuda M, Nakajima N, Hiroshima K, Inadera H, Matsushima K, Kuriyama T. Plasma monocyte chemoattractant protein-1 and pulmonary vascular resistance in chronic thromboembolic pulmonary hypertension. *Am J Respir Crit Care Med*. 164:319-324, 2001.
353. Kurashima K, Fujimura M, Myou S, Kasahara K, Tachibana H, Amemiya N, Ishiura Y, Onai N, Matsushima K, Nakao S. Effects of oral steroids on blood CXCR3+ and CCR4+ T cells in patients with bronchial asthma. *Am J Respir Crit Care Med*. 164:754-758, 2001.

354. Misu T, Onodera H, Fujihara K, Matsushima K, Yoshie O, Okita N, Takase S, Itoyama Y. Chemokine receptor expression on T cells in blood and cerebrospinal fluid at relapse and remission of multiple sclerosis: imbalance of Th1/Th2-associated chemokine signaling. *J Neuroimmunol.* 114:207-212, 2001.
355. Mukai T, Iwasaki M, Gao P, Tomura M, Yashiro-Ohtani Y, Ono S, Murai M, Matsushima K, Kurimoto M, Kogo M, Matsuya T, Fujiwara H, Hamaoka T. IL-12 plays a pivotal role in LFA-1-mediated T cell adhesiveness by up-regulation of CCR5 expression. *J Leukoc Biol.* 70:422-430, 2001.
356. Nagai S, Hashimoto S, Yamashita T, Toyoda N, Satoh T, Suzuki T, Matsushima K. Comprehensive gene expression profile of human activated T(h)1- and T(h)2-polarized cells. *Int Immunol.* 13:367-376, 2001.
357. Nagase H, Kudo K, Izumi S, Ohta K, Kobayashi N, Yamaguchi M, Matsushima K, Morita Y, Yamamoto K, Hirai K. Chemokine receptor expression profile of eosinophils at inflamed tissue sites: Decreased CCR3 and increased CXCR4 expression by lung eosinophils. *J Allergy Clin Immunol.* 108:563-569, 2001.
358. Saji H, Koike M, Yamori T, Saji S, Seiki M, Matsushima K, Toi M. Significant correlation of monocyte chemoattractant protein-1 expression with neovascularization and progression of breast carcinoma. *Cancer.* 92:1085-1091, 2001.
359. Takaoka A, Tanaka Y, Tsuji T, Jinushi T, Hoshino A, Asakura Y, Mita Y, Watanabe K, Nakaike S, Togashi Y, Koda T, Matsushima K, Nishimura T. A critical role for mouse CXC chemokine(s) in pulmonary neutrophilia during Th type 1-dependent airway inflammation. *J Immunol.* 167:2349-2353, 2001.
360. Takayama T, Morelli AE, Onai N, Hirao M, Matsushima K, Tahara H, Thomson AW. Mammalian and viral IL-10 enhance C-C chemokine receptor 5 but down-regulate C-C chemokine receptor 7 expression by myeloid dendritic cells: impact on chemotactic responses and in vivo homing ability. *J Immunol.* 166:7136-7143, 2001.
361. Vestergaard C, Kirstejn N, Gesser B, Mortensen JT, Matsushima K, Larsen CG. IL-10 augments the IFN-gamma and TNF-alpha induced TARC production in HaCaT cells: a possible mechanism in the inflammatory reaction of atopic dermatitis. *J Dermatol Sci.* 26:46-54, 2001.
362. Wada T, Furuichi K, Sakai N, Hisada Y, Kobayashi K, Mukaida N, Tomosugi N, Matsushima K, Yokoyama H. Involvement of p38 mitogen-activated protein kinase followed by chemokine expression in crescentic glomerulonephritis. *Am J Kidney Dis.* 38:1169-1177, 2001.
363. Wada T, Yokoyama H, Matsushima K, Kobayashi K. Chemokines in renal diseases. *Int Immunopharmacol.* 1:637-645, 2001.
364. Wakugawa M, Nakamura K, Kakinuma T, Onai N, Matsushima K, Tamaki K. CC chemokine receptor 4 expression on peripheral blood CD4+ T cells reflects disease activity of atopic dermatitis. *J Invest Dermatol.* 117:188-196, 2001.
365. Watanabe Y, Hashizume M, Kataoka S, Hamaguchi E, Morimoto N, Tsuru S, Katoh S, Miyake K, Matsushima K, Tominaga M, Kurashige T, Fujimoto S, Kincade PW, Tominaga A.

- Differentiation stages of eosinophils characterized by hyaluronic acid binding via CD44 and responsiveness to stimuli. *DNA Cell Biol.* 20:189-202, 2001.
366. Yamashita T, Kaneko S, Hashimoto S, Sato T, Nagai S, Toyoda N, Suzuki T, Kobayashi K, Matsushima K. Serial analysis of gene expression in chronic hepatitis C and hepatocellular carcinoma. *Biochem Biophys Res Commun.* 282:647-654, 2001.
367. Yeh M, Leitinger N, de Martin R, Onai N, Matsushima K, Vora DK, Berliner JA, Reddy ST. Increased transcription of IL-8 in endothelial cells is differentially regulated by TNF-alpha and oxidized phospholipids. *Arterioscler Thromb Vasc Biol.* 21:1585-1591, 2001.
368. Yokochi S, Hashimoto H, Ishiwata Y, Shimokawa H, Haino M, Terashima Y, Matsushima K. An anti-inflammatory drug, propagermanium, may target GPI-anchored proteins associated with an MCP-1 receptor, CCR2. *J Interferon Cytokine Res.* 21:389-398, 2001.
369. Yoneyama H, Matsuno K, Zhang Y, Murai M, Itakura M, Ishikawa S, Hasegawa G, Naito M, Asakura H, Matsushima K. Regulation by chemokines of circulating dendritic cell precursors, and the formation of portal tract-associated lymphoid tissue, in a granulomatous liver disease. *J Exp Med.* 193:35-49, 2001.
370. Dong H, Toyoda N, Yoneyama H, Kurachi M, Kasahara T, Kobayashi Y, Inadera H, Hashimoto S, Matsushima K. Gene expression profile analysis of the mouse liver during bacteria-induced fulminant hepatitis by a cDNA microarray system. *Biochem Biophys Res Commun.* 298:675-686, 2002.
371. Furuichi K, Wada T, Iwata Y, Sakai N, Yoshimoto K, Kobayashi Ki K, Mukaida N, Matsushima K, Yokoyama H. Administration of FR167653, a new anti-inflammatory compound, prevents renal ischaemia/reperfusion injury in mice. *Nephrol Dial Transplant.* 17:399-407, 2002.
372. Harada K, Tsuneyama K, Yasoshima M, Kanemori Y, Ohta H, Masuda S, Onai N, Matsushima K, Nakanuma Y. Type1 and type2 memory T cells imbalance shown by expression of intrahepatic chemokine receptors relates to pathogenesis of primary biliary cirrhosis. *Hepatol Res.* 24:290, 2002.
373. Higashi N, Fujioka K, Denda-Nagai K, Hashimoto S, Nagai S, Sato T, Fujita Y, Morikawa A, Tsuiji M, Miyata-Takeuchi M, Sano Y, Suzuki N, Yamamoto K, Matsushima K, Irimura T. The macrophage C-type lectin specific for galactose/N-acetylgalactosamine is an endocytic receptor expressed on monocyte-derived immature dendritic cells. *J Biol Chem.* 277:20686-20693, 2002.
374. Inadera H, Dong HY, Matsushima K. WISP-2 is a secreted protein and can be a marker of estrogen exposure in MCF-7 cells. *Biochem Biophys Res Commun.* 294:602-608, 2002.
375. Inadera H, Nagai S, Dong HY, Matsushima K. Molecular analysis of lipid-depleting factor in a colon-26-inoculated cancer cachexia model. *Int J Cancer.* 101:37-45, 2002.
376. Ishikawa S, Nagai S, Sato T, Akadegawa K, Yoneyama H, Zhang YY, Onai N, Matsushima K. Increased circulating CD11b+CD11c+ dendritic cells (DC) in aged BWF1 mice which can be matured by TNF-alpha into BLC/CXCL13-producing DC. *Eur J Immunol.* 32:1881-1887, 2002.
377. Kuno K, Baba C, Asaka A, Matsushima C, Matsushima K, Hosono R. The *Caenorhabditis*

- elegans ADAMTS family gene *adt-1* is necessary for morphogenesis of the male copulatory organs. *J Biol Chem.* 277:12228-12236, 2002.
378. Kurachi M, Hashimoto S, Obata A, Nagai S, Nagahata T, Inadera H, Sone H, Tohyama C, Kaneko S, Kobayashi K, Matsushima K. Identification of 2,3,7,8-tetrachlorodibenzo-p-dioxin-responsive genes in mouse liver by serial analysis of gene expression. *Biochem Biophys Res Commun.* 292:368-377, 2002.
379. Masuda M, Senju S, Fujii Si S, Terasaki Y, Takeya M, Hashimoto Si S, Matsushima K, Yumoto E, Nishimura Y. Identification and immunocytochemical analysis of DCNP1, a dendritic cell-associated nuclear protein. *Biochem Biophys Res Commun.* 290:1022-1029, 2002.
380. Nagase H, Miyamasu M, Yamaguchi M, Imanishi M, Tsuno NH, Matsushima K, Yamamoto K, Morita Y, Hirai K. Cytokine-mediated regulation of CXCR4 expression in human neutrophils. *J Leukoc Biol.* 71:711-717, 2002.
381. Narumi S, Kaburaki T, Yoneyama H, Iwamura H, Kobayashi Y, Matsushima K. Neutralization of IFN-inducible protein 10/CXCL10 exacerbates experimental autoimmune encephalomyelitis. *Eur J Immunol.* 32:1784-1791, 2002.
382. Obata-Onai A, Hashimoto S, Onai N, Kurachi M, Nagai S, Shizuno K, Nagahata T, Matsushima K. Comprehensive gene expression analysis of human NK cells and CD8+ T lymphocytes. *Int Immunol.* 14(10):1085-1098, 2002
383. Okazaki H, Kakurai M, Hirata D, Sato H, Kamimura T, Onai N, Matsushima K, Nakagawa H, Kano S, Minota S. Characterization of chemokine receptor expression and cytokine production in circulating CD4+ T cells from patients with atopic dermatitis: up-regulation of C-C chemokine receptor 4 in atopic dermatitis. *Clin Exp Allergy.* 32:1236-1242, 2002.
384. Onai N, Kitabatake M, Zhang YY, Ishikawa H, Ishikawa S, Matsushima K. Pivotal role of CCL25 (TECK)-CCR9 in the formation of gut cryptopatches and consequent appearance of intestinal intraepithelial T lymphocytes. *Int Immunol.* 14:687-694, 2002.
385. Osman MO, Gesser B, Mortensen JT, Matsushima K, Jensen SL, Larsen CG. Profiles of pro-inflammatory cytokines in the serum of rabbits after experimentally induced acute pancreatitis. *Cytokine.* 17:53-59, 2002.
386. Sakai N, Wada T, Furuichi K, Iwata Y, Yoshimoto K, Kitagawa K, Kokubo S, Kobayashi M, Takeda S, Kida H, Kobayashi K, Mukaida N, Matsushima K, Yokoyama H. p38 MAPK phosphorylation and NF-kappa B activation in human crescentic glomerulonephritis. *Nephrol Dial Transplant.* 17:998-1004, 2002.
387. Sekiya T, Yamada H, Yamaguchi M, Yamamoto K, Ishii A, Yoshie O, Sano Y, Morita A, Matsushima K, Hirai K. Increased levels of a TH2-type CC chemokine thymus and activation-regulated chemokine (TARC) in serum and induced sputum of asthmatics. *Allergy.* 57:173-177, 2002.
388. Tozawa Y, Matsushima K. Peripheral 5-HT(2A)-receptor-mediated formation of an inhibitor of atrial natriuretic peptide binding involves inflammation. *Eur J Pharmacol.* 440:37-44, 2002.
389. Tsuchihashi Y, Oishi K, Yoshimine H, Suzuki S, Kumatori A, Sunazuka T, Omura S,

- Matsushima K, Nagatake T. Fourteen-member macrolides suppress interleukin-8 production but do not promote apoptosis of activated neutrophils. *Antimicrob Agents Chemother.* 46:1101-1104, 2002.
390. Tsuda H, Michimata T, Hayakawa S, Tanebe K, Sakai M, Fujimura M, Matsushima K, Saito S. A Th2 chemokine, TARC, produced by trophoblasts and endometrial gland cells, regulates the infiltration of CCR4+ T lymphocytes into human decidua at early pregnancy. *Am J Reprod Immunol.* 48:1-8, 2002.
391. Uekusa Y, Yu WG, Mukai T, Gao P, Yamaguchi N, Murai M, Matsushima K, Obika S, Imanishi T, Higashibata Y, Nomura S, Kitamura Y, Fujiwara H, Hamaoka T. A pivotal role for CC chemokine receptor 5 in T-cell migration to tumor sites induced by interleukin 12 treatment in tumor-bearing mice. *Cancer Res.* 62:3751-3758, 2002.
392. Ueno T, Hara K, Willis MS, Malin MA, Hopken UE, Gray DH, Matsushima K, Lipp M, Springer TA, Boyd RL, Yoshie O, Takahama Y. Role for CCR7 ligands in the emigration of newly generated T lymphocytes from the neonatal thymus. *Immunity.* 16:205-218, 2002.
393. Wang HY, Matsui M, Araya S, Onai N, Matsushima K, Saida T. Chemokine receptors associated with immunity within and outside the central nervous system in early relapsing-remitting multiple sclerosis. *J Neuroimmunol.* 133:184-192, 2002.
394. Wang Y, Zhang Y, Yoneyama H, Onai N, Sato T, Matsushima K. Identification of CD8alpha+CD11c- lineage phenotype-negative cells in the spleen as committed precursor of CD8alpha+ dendritic cells. *Blood.* 100:569-577, 2002.
395. Yokoyama H, Wada T, Kobayashi K, Kuno K, Kurihara H, Shindo T, Matsushima K. A disintegrin and metalloproteinase with thrombospondin motifs (ADAMTS)-1 null mutant mice develop renal lesions mimicking obstructive nephropathy. *Nephrol Dial Transplant.* 17 Suppl 9:39-41, 2002.
396. Yoneyama H, Narumi S, Zhang Y, Murai M, Baggiolini M, Lanzavecchia A, Ichida T, Asakura H, Matsushima K. Pivotal role of dendritic cell-derived CXCL10 in the retention of T helper cell 1 lymphocytes in secondary lymph nodes. *J Exp Med.* 195:1257-1266, 2002.
397. Yoshie O, Fujisawa R, Nakayama T, Harasawa H, Tago H, Izawa D, Hieshima K, Tatsumi Y, Matsushima K, Hasegawa H, Kanamaru A, Kamihira S, Yamada Y. Frequent expression of CCR4 in adult T-cell leukemia and human T-cell leukemia virus type 1-transformed T cells. *Blood.* 99:1505-1511, 2002.
398. Hashimoto SI, Nagai S, Sese J, Suzuki T, Obata A, Sato T, Toyoda N, Dong HY, Kurachi M, Nagahata T, Shizuno KI, Morishita S, Matsushima K. Gene expression profile in human leukocytes. *Blood.* 101:3509-3513, 2003.
399. Iwata Y, Wada T, Furuichi K, Sakai N, Matsushima K, Yokoyama H, Kobayashi K. p38 Mitogen-Activated Protein Kinase Contributes to Autoimmune Renal Injury in MRL-Fas(lpr) Mice. *J Am Soc Nephrol.* 14:57-67, 2003.
400. Kakinuma T, Sugaya M, Nakamura K, Kaneko F, Wakugawa M, Matsushima K, Tamaki K. Thymus and activation-regulated chemokine (TARC/CCL17) in mycosis fungoides: Serum TARC levels reflect the disease activity of mycosis fungoides. *J Am Acad Dermatol.* 48:23-30, 2003.

401. Murai M, Yoneyama H, Ezaki T, Suematsu M, Terashima Y, Harada A, Hamada H, Asakura H, Ishikawa H, Matsushima K. Peyer's patch is the essential site in initiating murine acute and lethal graft-versus-host reaction. *Nat Immunol.* 4:154-160, 2003.
402. Schaidler H, Oka M, Bogenrieder T, Nesbit M, Satyamoorthy K, Berking C, Matsushima K, Herlyn M. Differential response of primary and metastatic melanomas to neutrophils attracted by IL-8. *Int J Cancer.* 103:335-343, 2003.
403. Yamaryo T, Oishi K, Yoshimine H, Tsuchihashi Y, Matsushima K, Nagatake T. Fourteen-member macrolides promote the phosphatidylserine receptor-dependent phagocytosis of apoptotic neutrophils by alveolar macrophages. *Antimicrob Agents Chemother.* 47:48-53, 2003.
404. Kakinuma T, Wakugawa M, Nakamura K, Hino H, Matsushima K, Tamaki K. High level of thymus and activation-regulated chemokine in blister fluid and sera of patients with bullous pemphigoid. *Br J Dermatol.* 148:203-210, 2003.
405. Wada T, Yokoyama H, Matsushima K, Kobayashi K. Monocyte chemoattractant protein-1: does it play a role in diabetic nephropathy? *Nephrol Dial Transplant.* 18:457-9, 2003.
406. Nakae S, Komiyama Y, Narumi S, Sudo K, Horai R, Tagawa YI, Sekikawa K, Matsushima K, Asano M, Iwakura Y. IL-1-induced tumor necrosis factor-alpha elicits inflammatory cell infiltration in the skin by inducing IFN-gamma-inducible protein 10 in the elicitation phase of the contact hypersensitivity response. *Int Immunol.* 15:251-260, 2003.
407. Takizawa H, Abe S, Okazaki H, Kohyama T, Sugawara I, Saito Y, Ohtoshi T, Kawasaki S, Desaki M, Nakahara K, Yamamoto K, Matsushima K, Tanaka M, Sagai M, Kudoh S. Diesel Exhaust Particles Upregulate Eotaxin Gene Expression in Human Bronchial Epithelial Cells via Nuclear Factor- κ B-Dependent Pathway. *Am J Physiol Lung Cell Mol Physiol.* 284:L1055-1062, 2003.
408. Sekiya T, Tsunemi Y, Miyamasu M, Ohta K, Morita A, Saeki H, Matsushima K, Yoshie O, Tsuchiya N, Yamaguchi M, Yamamoto K, Tamaki K, Hirai K. Variations in the human Th2-specific chemokine TARC gene. *Immunogenetics.* 54:742-745, 2003.
409. Sasaki S, Yoneyama H, Suzuki K, Suriki H, Aiba T, Watanabe S, Kawauchi Y, Kawachi H, Shimizu F, Matsushima K, Asakura H, Narumi S. Blockade of CXCL10 protects mice from acute colitis and enhances crypt cell survival. *Eur J Immunol.* 32:3197-3205, 2002.
410. Eto Y, Shimokawa H, Tanaka E, Morishige K, Fuchigami M, Ishiwata Y, Matsushima K, Takeshita A. Long-term Treatment with Propagermanium Suppresses Atherosclerosis in WHHL Rabbits. *J Cardiovasc Pharmacol.* 41:171-177, 2003.
411. Shimokawa H, Eto Y, Miyata K, Morishige K, Kandabashi T, Matsushima K, Takeshita A. Propagermanium Suppresses Macrophage-Mediated Formation of Coronary Arteriosclerotic Lesions in Pigs in Vivo. *J Cardiovasc Pharmacol.* 41:372-380, 2003.
412. Yoshimura-Uchiyama C, Yamaguchi M, Nagase H, Fujisawa T, Ra C, Matsushima K, Iwata T, Igarashi T, Yamamoto K, Hirai K. Comparative effects of basophil-directed growth factors. *Biochem Biophys Res Commun.* 302:201-206, 2003.
413. Furuichi K, Wada T, Iwata Y, Kitagawa K, Kobayashi K, Hashimoto H, Ishiwata Y, Tomosugi N, Mukaida N, Matsushima K, Egashira K, Yokoyama H. Gene therapy expressing

- amino-terminal truncated monocyte chemoattractant protein-1 prevents renal ischemia-reperfusion injury. *J Am Soc Nephrol.* 14:1066-1071, 2003.
414. Jo Y, Matsumoto T, Yada S, Fujisawa K, Esaki M, Onai N, Matsushima K, Iida M. CCR4 is an up-regulated chemokine receptor of peripheral blood memory CD4+ T cells in Crohn's disease. *Clin Exp Immunol.* 132:332-338, 2003.
415. Kaburaki T, Fujino Y, Kawashima H, Merino G, Numaga J, Chen J, Matsushima K. Plasma and whole-blood chemokine levels in patients with Behcet's disease. *Graefes Arch Clin Exp Ophthalmol.* 241:353-358, 2003.
416. Katoh S, Fukushima K, Matsumoto N, Matsumoto K, Abe K, Onai N, Matsushima K, Matsukura S. Accumulation of CCR4-expressing CD4+ T cells and high concentration of its ligands (TARC and MDC) in bronchoalveolar lavage fluid of patients with eosinophilic pneumonia. *Allergy.* 58:518-523, 2003.
417. Toyoda N, Nagai S, Terashima Y, Motomura K, Haino M, Hashimoto S, Takizawa H, Matsushima K. Analysis of mRNA with microsomal fractionation using a SAGE-based DNA microarray system facilitates identification of the genes encoding secretory proteins. *Genome Res.* 13:1728-36, 2003.
418. Yoshimura-Uchiyama C, Yamaguchi M, Nagase H, Matsushima K, Igarashi T, Iwata T, Yamamoto K, Hirai K. Changing expression of IL-3 and IL-5 receptors in cultured human eosinophils. *Biochem Biophys Res Commun.* 309:26-31, 2003.
419. Komiya A, Nagase H, Yamada H, Sekiya T, Yamaguchi M, Sano Y, Hanai N, Furuya A, Ohta K, Matsushima K, Yoshie O, Yamamoto K, Hirai K. Concerted expression of eotaxin-1, eotaxin-2, and eotaxin-3 in human bronchial epithelial cells. *Cell Immunol.* 225:91-100, 2003.
420. Hashimoto S, Morohoshi K, Suzuki T, Matsushima K. Lipopolysaccharide-inducible gene expression profile in human monocytes. *Scand J Infect Dis.* 35:619-27, 2003.
421. Matsui M, Araya S, Wang HY, Onai N, Matsushima K, Saida T. Circulating lymphocyte subsets linked to intracellular cytokine profiles in normal humans. *Clin Exp Immunol.* 134:225-31, 2003.
422. Wang HY, Matsui M, Araya S, Onai N, Matsushima K, Saida T. Immune parameters associated with early treatment effects of high-dose intravenous methylprednisolone in multiple sclerosis. *J Neurol Sci.* 216:61-6, 2003.
423. Nagase H, Okugawa S, Ota Y, Yamaguchi M, Tomizawa H, Matsushima K, Ohta K, Yamamoto K, Hirai K. Expression and function of Toll-like receptors in eosinophils: activation by Toll-like receptor 7 ligand. *J Immunol.* 171:3977-82, 2003.
424. Furuichi K, Wada T, Iwata Y, Kitagawa K, Kobayashi K, Hashimoto H, Ishiwata Y, Asano M, Wang H, Matsushima K, Takeya M, Kuziel WA, Mukaida N, Yokoyama H. CCR2 Signaling Contributes to Ischemia-Reperfusion Injury in Kidney. *J Am Soc Nephrol.* 14:2503-15, 2003.
425. Ishida T, Utsunomiya A, Iida S, Inagaki H, Takatsuka Y, Kusumoto S, Takeuchi G, Shimizu S, Ito M, Komatsu H, Wakita A, Eimoto T, Matsushima K, Ueda R. Clinical significance of CCR4 expression in adult T-cell leukemia/lymphoma: its close association with skin involvement and unfavorable outcome. *Clin Cancer Res.* 9(10 Pt 1):3625-34, 2003.
426. Zhang Y, Yoneyama H, Wang Y, Ishikawa S, Hashimoto SI, Gao JL, Murphy P, Matsushima K.

- Mobilization of Dendritic Cell Precursors Into the Circulation by Administration of MIP-1 α in Mice. *J Natl Cancer Inst.* 96:201-9, 2004.
427. Ito T, Ishikawa S, Sato T, Akadegawa K, Yurino H, Kitabatake M, Hontsu S, Ezaki T, Kimura H, Matsushima K. Defective B1 cell homing to the peritoneal cavity and preferential recruitment of B1 cells in the target organs in a murine model for SLE. *J Immunol.* 172:3628-34, 2004.
428. Yoneyama H, Matsuno K, Zhang Y, Nishiwaki T, Kitabatake M, Ueha S, Narumi S, Morikawa S, Ezaki T, Lu B, Gerard C, Ishikawa S, Matsushima K. Evidence for recruitment of plasmacytoid dendritic cell precursors to inflamed lymph nodes through high endothelial venules. *Int Immunol.* 16:915-28, 2004.
429. Nishiwaki T, Yoneyama H, Eishi Y, Matsuo N, Tatsumi K, Kimura H, Kuriyama T, Matsushima K. Indigenous Pulmonary *Propionibacterium acnes* Primes the Host in the Development of Sarcoid-like Pulmonary Granulomatosis in Mice. *Am J Pathol.* 165:631-9, 2004.
430. Sato T, Ishikawa S, Akadegawa K, Ito T, Yurino H, Kitabatake M, Yoneyama H, Matsushima K. Aberrant B1 cell migration into the thymus results in activation of CD4 T cells through its potent antigen presenting activity in the development of murine lupus. *Eur J Immunol.* 34:3346-58, 2004.
431. Hontsu S, Yoneyama H, Ueha S, Terashima Y, Kitabatake M, Nakano A, Ito T, Kimura H, Matsushima K. Visualization of naturally occurring Foxp3+ regulatory T cells in normal and tumor-bearing mice. *Int Immunopharmacol.* 4:1785-93, 2004.
432. Hashimoto S, Suzuki Y, Kasai Y, Morohoshi K, Yamada T, Sese J, Morishita S, Sugano S, Matsushima K. 5'-end SAGE for the analysis of transcriptional start sites. *Nat Biotechnol.* 22:1146-9, 2004.
433. Wada T, Furuichi K, Sakai N, Iwata Y, Kitagawa K, Ishida Y, Kondo T, Hashimoto H, Ishiwata Y, Mukaida N, Tomosugi N, Matsushima K, Egashira K, Yokoyama H. Gene therapy via blockade of monocyte chemoattractant protein-1 for renal fibrosis. *J Am Soc Nephrol.* 15:940-8, 2004.
434. Kamohara H, Matsuyama W, Shimozato O, Abe K, Galligan C, Hashimoto S, Matsushima K, Yoshimura T. Regulation of tumour necrosis factor-related apoptosis-inducing ligand (TRAIL) and TRAIL receptor expression in human neutrophils. *Immunology.* 111:186-94, 2004.
435. Niwa R, Shoji-Hosaka E, Sakurada M, Shinkawa T, Uchida K, Nakamura K, Matsushima K, Ueda R, Hanai N, Shitara K. Defucosylated chimeric anti-CC chemokine receptor 4 IgG1 with enhanced antibody-dependent cellular cytotoxicity shows potent therapeutic activity to T-cell leukemia and lymphoma. *Cancer Res.* 64:2127-33, 2004.
436. Matsui M, Araya SI, Wang HY, Matsushima K, Saida T. Immunomonitoring measures in relapsing-remitting multiple sclerosis. *J Neuroimmunol.* 148:192-9, 2004.
437. Kotani M, Kotani T, Ishizaka A, Fujishima S, Koh H, Tasaka S, Sawafuji M, Ikeda E, Moriyama K, Kotake Y, Morisaki H, Aikawa N, Ohashi A, Matsushima K, Huang YC, Takeda J. Neutrophil depletion attenuates interleukin-8 production in mild-overstretch ventilated normal rabbit lung. *Crit Care Med.* 32:514-9, 2004.
438. Yoshimura-Uchiyama C, Iikura M, Yamaguchi M, Nagase H, Ishii A, Matsushima K,

- Yamamoto K, Shichijo M, Bacon KB, Hirai K. Differential modulation of human basophil functions through prostaglandin D2 receptors DP and chemoattractant receptor-homologous molecule expressed on Th2 cells/DP2. *Clin Exp Allergy*. 34:1283-90, 2004.
439. Inoue T, Fujishima S, Ikeda E, Yoshie O, Tsukamoto N, Aiso S, Aikawa N, Kubo A, Matsushima K, Yamaguchi K. CCL22 and CCL17 in rat radiation pneumonitis and in human idiopathic pulmonary fibrosis. *Eur Respir J*. 24:49-56, 2004.
440. Kitagawa K, Wada T, Furuichi K, Hashimoto H, Ishiwata Y, Asano M, Takeya M, Kuziel WA, Matsushima K, Mukaida N, Yokoyama H. Blockade of CCR2 ameliorates progressive fibrosis in kidney. *Am J Pathol*. 165:237-46, 2004.
441. Kuno K, Bannai K, Hakozaki M, Matsushima K, Hirose K. The carboxyl-terminal half region of ADAMTS-1 suppresses both tumorigenicity and experimental tumor metastatic potential. *Biochem Biophys Res Commun*. 319:1327-33, 2004.
442. Katoh S, Matsumoto N, Matsumoto K, Tokojima M, Ashitani J, Nakamura-Uchiyama F, Matsushima K, Matsukura S, Nawa Y. A possible role of TARC in antigen-specific Th2-dominant responses in patients with *Paragonimiasis westermani*. *Int Arch Allergy Immunol*. 134:248-52, 2004.
443. Yurino H, Ishikawa S, Sato T, Akadegawa K, Ito T, Ueha S, Inadera H, Matsushima K. Endocrine disruptors (environmental estrogens) enhance autoantibody production by B1 cells. *Toxicol Sci*. 81:139-47, 2004.
444. Fujimura T, Aiba S, Yoshino Y, Kuroki S, Kimura Y, Kikuchi K, Kunii T, Matsunaga J, Matsushima K, Tagami H. CCR4 Expression by atypical T cells in systemic pilotropic lymphoma: its behavior under treatment with interferon gamma, topical PUVA and systemic retinoid. *Dermatology*. 208:221-6, 2004.
445. Morimoto J, Yoneyama H, Shimada A, Shigihara T, Yamada S, Oikawa Y, Matsushima K, Saruta T, Narumi S. CXC chemokine ligand 10 neutralization suppresses the occurrence of diabetes in nonobese diabetic mice through enhanced beta cell proliferation without affecting insulinitis. *J Immunol*. 173:7017-24, 2004.
446. Motomura K, Toyoda N, Oishi K, Sato H, Nagai S, Hashimoto S, Tugume SB, Enzama R, Mugewa R, Mutuluza CK, Mugyeyi P, Nagatake T, Matsushima K. Identification of a host gene subset related to disease prognosis of HIV-1 infected individuals. *Int Immunopharmacol*. 4:1829-36, 2004.
447. Yanaba K, Mukaida N, Matsushima K, Murphy PM, Takehara K, Sato S. Role of C-C chemokine receptors 1 and 5 and CCL3/macrophage inflammatory protein-1alpha in the cutaneous Arthus reaction: possible attenuation of their inhibitory effects by compensatory chemokine production. *Eur J Immunol*. 34:3553-61, 2005.
448. Matsui M, Araya SI, Wang HY, Matsushima K, Saida T. Differences in systemic and central nervous system cellular immunity relevant to relapsing-remitting multiple sclerosis. *J Neurol*. 252:908-15, 2005.
449. Iizasa H, Yoneyama H, Mukaida N, Katakoka Y, Naito M, Yoshida N, Nakashima E, Matsushima K. Exacerbation of granuloma formation in IL-1 receptor antagonist-deficient mice with impaired dendritic cell maturation associated with Th2 cytokine production. *J*

- Immunol. 174:3273-80, 2005.
450. Matsuzaki Y, Hashimoto S, Fujita T, Suzuki T, Sakurai T, Matsushima K, Kawakami Y. Systematic identification of human melanoma antigens using serial analysis of gene expression (SAGE). *J Immunother.* 28:10-9, 2005.
451. Kasai Y, Hashimoto S, Yamada T, Sese J, Sugano S, Matsushima K, Morishita S. 5'SAGE: 5'-end Serial Analysis of Gene Expression database. *Nucleic Acids Res.* 33(Database issue):D550-2, 2005.
452. Ohta M, Tateishi K, Kanai F, Ueha S, Guleng B, Washida M, Tanaka Y, Ijichi H, Ikenoue T, Sata M, Sudo T, Shiina S, Kawabe T, Matsushima K, Omata M. Reduced p38 mitogen-activated protein kinase in donor grafts accelerates acute intestinal graft-versus-host disease in mice. *Eur J Immunol.* 35:2210-21, 2005.
453. Akadegawa K, Ishikawa S, Sato T, Suzuki J, Yurino H, Kitabatake M, Ito T, Kuriyama T, Matsushima K. Breakdown of mucosal immunity in the gut and resultant systemic sensitization by oral antigens in a murine model for systemic lupus erythematosus. *J Immunol.* 174:5499-506, 2005.
454. Niwa R, Sakurada M, Kobayashi Y, Uehara A, Matsushima K, Ueda R, Nakamura K, Shitara K. Enhanced natural killer cell binding and activation by low-fucose IgG1 antibody results in potent antibody-dependent cellular cytotoxicity induction at lower antigen density. *Clin Cancer Res.* 11:2327-36, 2005.
455. Takeuchi A, Usui Y, Takeuchi M, Hattori T, Kezuka T, Suzuki J, Okunuki Y, Iwasaki T, Haino M, Matsushima K, Usui M. CCR5-Deficient Mice Develop Experimental Autoimmune Uveoretinitis in the Context of a Deviant Effector Response. *Invest Ophthalmol Vis Sci.* 46:3753-60, 2005.
456. Iikura M, Suzukawa M, Yamaguchi M, Sekiya T, Komiya A, Yoshimura-Uchiyama C, Nagase H, Matsushima K, Yamamoto K, Hirai K. 5-Lipoxygenase products regulate basophil functions: 5-Oxo-ETE elicits migration, and leukotriene B(4) induces degranulation. *J Allergy Clin Immunol.* 116:578-85, 2005.
457. Yoneyama H, Matsuno K, Toda E, Nishiwaki T, Matsuo N, Nakano A, Narumi S, Lu B, Gerard C, Ishikawa S, Matsushima K. Plasmacytoid DCs help lymph node DCs to induce anti-HSV CTLs. *J Exp Med.* 202:425-35, 2005.
458. Tokuyama H, Ueha S, Kurachi M, Matsushima K, Moriyasu F, Blumberg RS, Kakimi K. The simultaneous blockade of chemokine receptors CCR2, CCR5 and CXCR3 by a non-peptide chemokine receptor antagonist protects mice from dextran sodium sulfate-mediated colitis. *Int Immunol.* 17:1023-34, 2005.
459. Terashima Y, Onai N, Murai M, Enomoto M, Poonpiriya V, Hamada T, Motomura K, Suwa M, Ezaki T, Haga T, Kanegasaki S, Matsushima K. Pivotal function for cytoplasmic protein FROUNT in CCR2-mediated monocyte chemotaxis. *Nat Immunol.* 6:827-35, 2005.
460. Shigihara T, Shimada A, Oikawa Y, Yoneyama H, Kanazawa Y, Okubo Y, Matsushima K, Yamato E, Miyazaki J, Kasuga A, Saruta T, Narumi S. CXCL10 DNA vaccination prevents spontaneous diabetes through enhanced beta cell proliferation in NOD mice. *J Immunol.* 175:8401-8, 2005.

461. Hashida N, Ohguro N, Nakai K, Kobashi-Hashida M, Hashimoto S, Matsushima K, Tano Y. Microarray analysis of cytokine and chemokine gene expression after prednisolone treatment in murine experimental autoimmune uveoretinitis. *Invest Ophthalmol Vis Sci.* 46:4224-348, 2005.
462. Shozu M, Minami N, Yokoyama H, Inoue M, Kurihara H, Matsushima K, Kuno K. ADAMTS-1 is involved in normal follicular development, ovulatory process and organization of the medullary vascular network in the ovary. *J Mol Endocrinol.* 35:343-558, 2005.
463. Ezaki T, Kuwahara K, Morikawa S, Shimizu K, Sakaguchi N, Matsushima K, Matsuno K. Production of two novel monoclonal antibodies that distinguish mouse lymphatic and blood vascular endothelial cells. *Anat Embryol (Berl).* 211:379-93, 2006.
464. Furuichi K, Wada T, Iwata Y, Kokubo S, Hara A, Yamahana J, Sugaya T, Iwakura Y, Matsushima K, Asano M, Yokoyama H, Kaneko S. Interleukin-1-dependent sequential chemokine expression and inflammatory cell infiltration in ischemia-reperfusion injury. *Crit Care Med* 34:2447-55, 2006.
465. Hara A, Wada T, Furuichi K, Sakai N, Kawachi H, Shimizu F, Shibuya M, Matsushima K, Yokoyama H, Egashira K, Kaneko S. Blockade of VEGF accelerates proteinuria, via decrease in nephrin expression in rat crescentic glomerulonephritis. *Kidney Int* 69:1986-95, 2006.
466. Hasan MR, Jin M, Matsushima K, Miyamoto S, Kotani S, Nakagawa H. Differences in the regulation of microtubule stability by the pro-rich region variants of microtubule-associated protein 4. *FEBS Lett* 580:3505-10, 2006.
467. Ishida Y, Kondo T, Kimura A, Matsushima K, Mukaida N. Absence of IL-1 receptor antagonist impaired wound healing along with aberrant NF-kappaB activation and a reciprocal suppression of TGF-beta signal pathway. *J Immunol* 176:5598-606, 2006.
468. Kanzawa N, Nishigaki K, Hayashi T, Ishii Y, Furukawa S, Niuro A, Yasui F, Kohara M, Morita K, Matsushima K, Le MQ, Masuda T, Kannagi M. Augmentation of chemokine production by severe acute respiratory syndrome coronavirus 3a/X1 and 7a/X4 proteins through NF-kappaB activation. *FEBS Lett.* 580:6807-12, 2006.
469. Komiya A, Nagase H, Okugawa S, Ota Y, Suzukawa M, Kawakami A, Sekiya T, Matsushima K, Ohta K, Hirai K, Yamamoto K, Yamaguchi M. Expression and function of toll-like receptors in human basophils. *Int Arch Allergy Immunol* 140 Suppl 1:23-7, 2006.
470. Kurashima K, Fujimura M, Myou S, Ishiura Y, Onai N, Matsushima K. Asthma severity is associated with an increase in both blood CXCR3+ and CCR4+ T cells. *Respirology* 11:152-7, 2006.
471. Kurita S, Koyama J, Onizuka S, Motomura K, Watanabe H, Watanabe K, Senba M, Apicella MA, Murphy TF, Yoneyama H, Matsushima K, Nagatake T, Oishi K. Dynamics of dendritic cell migration and the subsequent induction of protective immunity in the lung after repeated airway challenges by nontypeable *Haemophilus influenzae* outer membrane protein. *Vaccine* 24:5896-903, 2006.
472. Ohmori K, Fukui F, Kiso M, Imai T, Yoshie O, Hasegawa H, Matsushima K, Kannagi R. Identification of cutaneous lymphocyte-associated antigen as sialyl 6-sulfo Lewis X, a selectin ligand expressed on a subset of skin-homing helper memory T cells. *Blood* 107:3197-204,

- 2006.
473. Oka M, Norose K, Matsushima K, Nishigori C, Herlyn M. Overexpression of IL-8 in the cornea induces ulcer formation in the SCID mouse. *Br J Ophthalmol* 90:612-5, 2006.
474. Sakai N, Wada T, Furuichi K, Shimizu K, Kokubo S, Hara A, Yamahana J, Okumura T, Matsushima K, Yokoyama H, Kaneko S. MCP-1/CCR2-dependent loop for fibrogenesis in human peripheral CD14-positive monocytes. *J Leukoc Biol* 79:555-63, 2006.
475. Sakai N, Wada T, Yokoyama H, Lipp M, Ueha S, Matsushima K, Kaneko S. Secondary lymphoid tissue chemokine (SLC/CCL21)/CCR7 signaling regulates fibrocytes in renal fibrosis. *Proc Natl Acad Sci U S A.* 103:14098-103, 2006.
476. Suzukawa M, Komiya A, Iikura M, Nagase H, Yoshimura-Uchiyama C, Yamada H, Kawasaki H, Ohta K, Matsushima K, Hirai K, Yamamoto K, Yamaguchi M. Trans-basement membrane migration of human basophils: role of matrix metalloproteinase-9. *nt Immunol.* 18:1575-83, 2006.
477. Tsunemi Y, Saeki H, Nakamura K, Nagakubo D, Nakayama T, Yoshie O, Kagami S, Shimazu K, Kadono T, Sugaya M, Komine M, Matsushima K, Tamaki K. CCL17 transgenic mice show an enhanced Th2-type response to both allergic and non-allergic stimuli. *Eur J Immunol* 36:2116-27, 2006.
478. Wada T, Azuma H, Furuichi K, Sakai N, Kitagawa K, Iwata Y, Matsushima K, Takahara S, Yokoyama H, Kaneko S. Reduction in Chronic Allograft Nephropathy by Inhibition of p38 Mitogen-Activated Protein Kinase. *Am J Nephrol* 26:319-25, 2006.
479. Zenclussen AC, Gerlof K, Zenclussen ML, Ritschel S, Zambon Bertoja A, Fest S, Hontsu S, Ueha S, Matsushima K, Leber J, Volk HD. Regulatory T cells induce a privileged tolerant microenvironment at the fetal-maternal interface. *Eur J Immunol* 36:82-94, 2006.
480. Zhu L, Ji F, Wang Y, Zhang Y, Liu Q, Zhang JZ, Matsushima K, Cao Q, Zhang Y. Synovial autoreactive T cells in rheumatoid arthritis resist IDO-mediated inhibition. *J Immunol.* 177:8226-33, 2006.
481. Ueha S, Murai M, Yoneyama H, Kitabatake M, Imai T, Shimaoka T, Yonehara S, Ishikawa S, Matsushima K. Intervention of MAdCAM-1 or Fractalkine alleviates graft-versus-host reaction associated intestinal injury while preserving graft-versus-tumor effects. *J Leukoc Biol.* 81:176-85, 2007.
482. Ishikawa S, Matsushima K. Aberrant B1 cell trafficking in a murine model for lupus. *Front Biosci.* 12:1790-803, 2007.
483. Kurachi M, Kakimi K, Ueha S, Matsushima K. Maintenance of memory CD8+ T cell diversity and proliferative potential by a primary response upon re-challenge. *Int Immunol.* 19:105-15, 2007.
484. Kitabatake M, Inoue S, Yasui F, Yokochi S, Arai M, Morita K, Shida H, Kidokoro M, Murai F, Le MQ, Mizuno K, Matsushima K, Kohara M. SARS-CoV spike protein-expressing recombinant vaccinia virus efficiently induces neutralizing antibodies in rabbits pre-immunized with vaccinia virus. *Vaccine.* 25:630-7, 2007.
485. Nakamoto Y, Mizukoshi E, Tsuji H, Sakai Y, Kitahara M, Arai K, Yamashita T, Yokoyama K, Mukaida N, Matsushima K, Matsui O, Kaneko S. Combined therapy of transcatheter hepatic

- arterial embolization with intratumoral dendritic cell infusion for hepatocellular carcinoma: clinical safety. *Clin Exp Immunol.* 147:296-305, 2007.
486. Vasilescu A, Terashima Y, Enomoto M, Heath S, Poonpiriya V, Gatanaga H, Do H, Diop G, Hirtzig T, Auewarakul P, Lauhakirti D, Sura T, Charneau P, Marullo S, Therwath A, Oka S, Kanegasaki S, Lathrop M, Matsushima K, Zagury J-F, Matsuda F. A haplotype of the human CXCR1 gene protective against rapid disease progression in HIV-1+ patients. *Proc Natl Acad Sci U S A.* 104:3354-9, 2007.
487. Ishida Y, Kimura A, Kondo T, Hayashi T, Ueno M, Takakura N, Matsushima K, Mukaida N. Essential roles of the CC chemokine ligand 3-CC chemokine receptor 5 axis in bleomycin-induced pulmonary fibrosis through regulation of macrophage and fibrocyte infiltration. *Am J Pathol.* 170:843-54, 2007.
488. Nakano A, Yoneyama H, Ueha S, Kitabatake M, Ishikawa S, Kawase I, Sugiyama H, Matsushima K. Intravenous administration of MIP-1alpha with intra-tumor injection of P. acnes shows potent anti-tumor effect. *Int Immunopharmacol.* 7:845-57, 2007.
489. Wada T, Sakai N, Matsushima K, Kaneko S. Fibrocytes: a new insight into kidney fibrosis. *Kidney Int.* 72:269-73, 2007.
490. Masahiro Kasahara, Kiyoshi Naruse, Shin Sasaki, Yoichiro Nakatani, Wei Qu, Budrul Ahsan, Tomoyuki Yamada, Yukinobu Nagayasu, Koichiro Doi, Yasuhiro Kasai, Tomoko Jindo, Daisuke Kobayashi, Atsuko Shimada, Atsushi Toyoda, Yoko Kuroki, Asao Fujiyama, Takashi Sasaki, Atsushi Shimizu, Shuichi Asakawa, Nobuyoshi Shimizu, Shin-ichi Hashimoto, Jun Yang, Yongjun Lee, Kouji Matsushima, Sumio Sugano, Mitsuru Sakaizumi, Takanori Narita, Kazuko Ohishi, Shinobu Haga, Fumiko Ohta, Hisayo Nomoto, Keiko Nogata, Tomomi Morishita, Tomoko Endo, Tadasu Shin-I, Hiroyuki Takeda, Shinichi Morishita and Yuji Kohara. The medaka draft genome: new insights into vertebrate genome evolution. *Nature.* 447:714-9, 2007.
491. Nakaya I, Wada T, Furuichi K, Sakai N, Ksitagawa K, Yokoyama H, Ishida Y, Kondo T, Sugaya T, Kawachi H, Shimizu F, Narumi S, Haino M, Gerard C, Matsushima K, Kaneko S. Blockade of IP-10/CXCR3 promotes progressive renal fibrosis. *Nephron Exp Nephrol.* 107:e12-21, 2007.
492. Ueha S, Yoneyama H, Hontsu S, Kurachi M, Kitabatake M, Abe J, Yoshie O, Shibayama S, Sugiyama T, Matsushima K. CCR7 mediates the migration of Foxp3+ regulatory T cells to the paracortical areas of peripheral lymph nodes through high endothelial venules. *J Leukoc Biol.* 82:1230-8, 2007.
493. Hao L, Zhang C, Qiu Y, Wang L, Luo Y, Jin M, Zhang Y, Guo TB, Matsushima K, Zhang Y. Recombination of CXCR4, VEGF, and MMP-9 predicting lymph node metastasis in human breast cancer. *Cancer Lett.* 253:34-42, 2007.
494. Kai Y, Yoneyama H, Koyama J, Hamada K, Kimura H, Matsushima K. Treatment with chondroitinase ABC alleviates bleomycin-induced pulmonary fibrosis. *Med Mol Morphol.* 40:128-40, 2007.
495. Shimaoka T, Seino K, Kume N, Minami M, Nishime C, Suematsu M, Kita T, Taniguchi M, Matsushima K, Yonehara S. Critical role for CXC chemokine ligand 16 (SR-PSOX) in Th1

- response mediated by NKT cells. *J Immunol.* 179:8172-9, 2007.
496. Fukushima K, Abiru N, Nagayama Y, Kobayashi M, Satoh T, Nakahara M, Kawasaki E, Yamasaki H, Ueha S, Matsushima K, Liu E, Eguchi K. Combined insulin B:9-23 self-peptide and polyinosinic-polycytidylic acid accelerate insulinitis but inhibit development of diabetes by increasing the proportion of CD4(+)Foxp3(+) regulatory T cells in the islets in non-obese diabetic mice. *Biochem Biophys Res Commun.* 367(4):719-24, 2008.
497. Guo Z, Jang MH, Otani K, Bai Z, Umemoto E, Matsumoto M, Nishiyama M, Yamasaki M, Ueha S, Matsushima K, Hirata T, Miyasaka M. CD4+CD25+ regulatory T cells in the small intestinal lamina propria show an effector/memory phenotype. *Int Immunol.* 20(3):307-15, 2008.
498. Ahsan B, Kobayashi D, Yamada T, Kasahara M, Sasaki S, Saito TL, Nagayasu Y, Doi K, Nakatani Y, Qu W, Jindo T, Shimada A, Naruse K, Toyoda A, Kuroki Y, Fujiyama A, Sasaki T, Shimizu A, Asakawa S, Shimizu N, Hashimoto S, Yang J, Lee Y, Matsushima K, Sugano S, Sakaizumi M, Narita T, Ohishi K, Haga S, Ohta F, Nomoto H, Nogata K, Morishita T, Endo T, Shin-I T, Takeda H, Kohara Y, Morishita S. UTGB/medaka: genomic resource database for medaka biology. *Nucleic Acids Res.* 36(Database issue):D747-52, 2008.
499. Shiraishi K, Ishiwata Y, Nakagawa K, Yokochi S, Taruki C, Akuta T, Ohtomo K, Matsushima K, Tamatani T, Kanegasaki S. Enhancement of antitumor radiation efficacy and consistent induction of the abscopal effect in mice by ECI301, an active variant of MIP-1 α . *Clin Cancer Res.* 14(4):1159-66, 2008.
500. Nanno M, Kanari Y, Naito T, Inoue N, Hisamatsu T, Chinen H, Sugimoto K, Shimomura Y, Yamagishi H, Shiohara T, Ueha S, Matsushima K, Suematsu M, Mizoguchi A, Hibi T, Bhan AK, Ishikawa H. Exacerbating role of gammadelta T cells in chronic colitis of T-cell receptor alpha mutant mice. *Gastroenterology.* 134:481-90, 2008.
501. Inadera H, Tachibana S, Takasaki I, Tabuchi Y, Matsushima K, Uchida M, Shimomura A. Expression profile of liver genes in response to hepatotoxicants identified using a SAGE-based customized DNA microarray system. *Toxicol Lett.* 177:20-30, 2008.
502. Sakai N, Wada T, Matsushima K, Bucala R, Iwai M, Horiuchi M, Kaneko S. The renin-angiotensin system contributes to renal fibrosis through regulation of fibrocytes. *J Hypertens.* 26:780-90, 2008.
503. Sawanobori Y, Ueha S, Kurachi M, Shimaoka T, Talmadge JE, Abe J, Shono Y, Kitabatake M, Kakimi K, Mukaida N, Matsushima K. Chemokine-mediated rapid turnover of myeloid-derived suppressor cells in tumor-bearing mice. *Blood.* 111:5457-66, 2008.
504. Wada T, Matsushima K, Kaneko S. The role of chemokines in glomerulonephritis. *Front Biosci.* 13:3966-74, 2008.
505. Furuichi K, Wada T, Kitajikma S, Toyama T, Okumura T, Hara A, Kawachi H, Shimizu F, Sugaya T, Mukaida N, Narumi S, Matsushima K, Kaneko S. IFN-inducible protein 10 (CXCL10) regulates tubular cell proliferation in renal ischemia-reperfusion injury. *Nephron Exp Nephrol.* 109:c29-38, 2008.
506. Shawkat S, Karima R, Tojo T, Tadakuma H, Saitoh S, Akashi-Takamura S, Miyake K, Funatsu T, Matsushima K. Visualization of the molecular dynamics of lipopolysaccharide on the

- plasma membrane of murine macrophages by total internal reflection fluorescence microscopy. *J Biol Chem.* 283:22962-71, 2008.
507. Lida N, Nakamoto Y, Baba T, Kakinoki K, Li YY, Wu Y, Matsushima K, Kaneko S, Mukaida N. Tumor cell apoptosis induces tumor-specific immunity in a CC chemokine receptor 1- and 5-dependent manner in mice. *J Leukoc Biol.* 84:1001-10, 2008.
508. Suzukawa M, Koketsu R, Iikura M, Nakae S, Matsumoto K, Nagase H, Saito H, Matsushima K, Ohta K, Yamamoto K, Yamaguchi M. Interleukin-33 enhances adhesion, CD11b expression and survival in human eosinophils. *Lab Invest.* 88:1245-53, 2008.
509. Komatsu K, Miyazaki D, Morohoshi K, Kuo CH, Kakimaru-Hasegawa A, Komatsu N, Namba S, Haino M, Matsushima K, Inoue Y. Pathogenesis of herpetic stromal keratitis in CCR5- and/or CXCR3-deficient mice. *Curr Eye Res.* 33:736-49, 2008.
510. Ahsan B, Saito TL, Hashimoto SI, Muramatsu K, Tsuda M, Sasaki A, Matsushima K, Aigaki T, Morishita S. MachiBase: a *Drosophila melanogaster* 5'-end mRNA transcription database. *Nucleic Acids Res.* 37(Database issue):D49-53, 2009.
511. Hashimoto S, Matsushima K. SAGE application in hematological research. *Curr Pharm Biotechnol.* 9(5):383-91. Review, 2008.
512. Suzukawa M, Iikura M, Koketsu R, Nagase H, Tamura C, Komiya A, Nakae S, Matsushima K, Ohta K, Yamamoto K, Yamaguchi M. An IL-1 cytokine member, IL-33, induces human basophil activation via its ST2 receptor. *J Immunol.* 181:5981-9, 2008.
513. Yasui F, Kai C, Kitabatake M, Inoue S, Yoneda M, Yokochi S, Kase R, Sekiguchi S, Morita K, Hishima T, Suzuki H, Karamatsu K, Yasutomi Y, Shida H, Kidokoro M, Mizuno K, Matsushima K, Kohara M. Prior immunization with severe acute respiratory syndrome (SARS)-associated coronavirus (SARS-CoV) nucleocapsid protein causes severe pneumonia in mice infected with SARS-CoV. *J Immunol.* 181:6337-48, 2008.
514. Wu Y, Li YY, Matsushima K, Baba T, Mukaida N. CCL3-CCR5 axis regulates intratumoral accumulation of leukocytes and fibroblasts and promotes angiogenesis in murine lung metastasis process. *J Immunol.* 181:6384-93, 2008.
515. Ishibe T, Kimura A, Ishida Y, Takayasu T, Hayashi T, Tsuneyama K, Matsushima K, Sakata I, Mukaida N, Kondo T. Reduced acetaminophen-induced liver injury in mice by genetic disruption of IL-1 receptor antagonist. *Lab Invest.* 89(1):68-79, 2009.
516. Yoshida A, Isomoto H, Hisatsune J, Nakayama M, Nakashima Y, Matsushima K, Mizuta Y, Hayashi T, Yamaoka Y, Azuma T, Moss J, Hirayama T, Kohno S. Enhanced expression of CCL20 in human *Helicobacter pylori*-associated gastritis. *Clin Immunol.* 130:290-7, 2009.
517. Abe J, Ueha S, Suzuki J, Tokano Y, Matsushima K, Ishikawa S. Increased Foxp3+ CD4+ Regulatory T Cells with Intact Suppressive Activity but Altered Cellular Localization in Murine Lupus. *Am J Pathol.* 173:1682-92, 2008.
518. Iwata Y, Furuichi K, Sakai N, Yamauchi H, Shinozaki Y, Zhou H, Kurokawa Y, Toyama T, Kitajima S, Okumura T, Yamada S, Maruyama I, Matsushima K, Kaneko S, Wada T. Dendritic Cells Contribute to Autoimmune Kidney Injury in MRL-Faslpr Mice. *J Rheumatol.* 36:306-14, 2009.
519. Hashimoto S, Qu W, Ahsan B, Ogoshi K, Sasaki A, Nakatani Y, Lee Y, Ogawa M, Ametani

- A, Suzuki Y, Sugano S, Lee CC, Nutter RC, Morishita S, Matsushima K. High-resolution analysis of the 5'-end transcriptome using a next generation DNA sequencer. *PLoS ONE*. 4:e4108, 2009.
520. Sasaki S, Mello CC, Shimada A, Nakatani Y, Hashimoto S, Ogawa M, Matsushima K, Gu SG, Kasahara M, Ahsan B, Sasaki A, Saito T, Suzuki Y, Sugano S, Kohara Y, Takeda H, Fire A, Morishita S. Chromatin-associated periodicity in genetic variation downstream of transcriptional start sites. *Science*. 323:401-4, 2009.
521. Grimstad O, Sawanobori Y, Vestergaard C, Bilsborough J, Olsen UB, Grønhøj-Larsen C, Matsushima K. Anti-interleukin-31-antibodies ameliorate scratching behaviour in NC/Nga mice: a model of atopic dermatitis. *Exp Dermatol*. 18:35-43, 2009.
522. International Silkworm Genome Consortium. The genome of a lepidopteran model insect, the silkworm *Bombyx mori*. *Insect Biochem Mol Biol*. 38:1036-45, 2008.
523. Tsuchihara K, Suzuki Y, Wakaguri H, Irie T, Tanimoto K, Hashimoto SI, Matsushima K, Mizushima-Sugano J, Yamashita R, Nakai K, Bentley D, Esumi H, Sugano S. Massive transcriptional start site analysis of human genes in hypoxia cells. *Nucleic Acids Res*. 37:2249-63, 2009.
524. Kumamoto M, Nishiwaki T, Matsuo N, Kimura H, Matsushima K. Minimally-cultured Bone Marrow mesenchymal stem cells ameliorate fibrotic lung injury. *Eur Respir J*. 34:740-8, 2009.
525. Ahsan B, Saito T L, Hashimoto S, Muramatsu K, Tsuda M, Sasaki A, Matsushima K, Aigaki T and Morishita S. MachiBase: a *Drosophila melanogaster* 5'-end mRNA transcription database. *Nucleic Acids Res*. 37(Database issue): D49-53, 2009.
526. Mizukoshi E, Nakamoto Y, Arai K, Yamashita T, Mukaida N, Matsushima K, Matsui O and Kaneko S. Enhancement of tumor-specific T cell responses by transcatheter arterial embolization with dendritic cell infusion for hepatocellular carcinoma. *Int J Cancer*. 126:2164-74, 2009.
527. Toda E, Terashima Y, Hirose K, Kanegasaki S, Matsushima K, FROUNT is a common regulator of CCR2 and CCR5 signaling to control directional migration. *Journal of Immunology*. 183:6387-94, 2009.
528. Kofuku Y, Yoshiura C, Ueda T, Terasawa H, Hirai T, Tominaga S, Hirose M, Maeda Y, Takahashi H, Terashima Y, Matsushima K, Shimada I. Structural basis of the interaction between chemokine stromal cell-derived factor-1/CXCL12 and its G-protein-coupled receptor CXCR4. *J Biol Chem*. 284:35240-50, 2009.
529. Komura T, Sakai Y, Honda M, Takamura T, Matsushima K, Kaneko S. CD14+monocytes are vulnerable and functionally impaired under ER stress in patients with type 2 diabetes. *Diabetes*. 59:634-43, 2009.
530. Hodo Y, Hashimoto SI, Honda M, Yamashita T, Suzuki Y, Sugano S, Kaneko S, Matsushima K. Comprehensive gene expression analysis of 5'-end of mRNA identified novel intronic transcripts associated with hepatocellular carcinoma. *Genomics*. 95:217-23, 2010.
531. Chamoto K, Wakita D, Ohkuri T, Uchinami Y, Matsushima K, Kitamura H, Nishimura T. 3-Methylcholanthrene-induced transforming growth factor-beta-producing carcinomas, but

- not sarcomas, are refractory to regulatory T cell-depletion therapy. *Cancer Sci.* 101:855-61, 2010.
532. Yamamoto K, Utsunomiya A, Tobinai K, Tsukasaki K, Uike N, Uozumi K, Yamaguchi K, Yamada Y, Hanada S, Tamura K, Nakamura S, Inagaki H, Ohshima K, Kiyoi H, Ishida T, Matsushima K, Akinaga S, Ogura M, Tomonaga M, Ueda R. Phase I study of KW-0761, a defucosylated humanized anti-CCR4 antibody, in relapsed patients with adult T-cell leukemia-lymphoma and peripheral T-cell lymphoma. *J Clin Oncol.* 28:1591-8, 2010.
533. Ikeuchi N, Futami J, Hosoi A, Noji S, Kurachi M, Ueha S, Fujii S, Yamada H, Matsushima K, Moriyasu F, Kakimi K. Efficient cross-presentation of soluble exogenous antigens introduced into dendritic cells using a weak-based amphiphilic peptide. *Biochem Biophys Res Commun.* 392:217-22, 2010.
534. Yoshiura C, Kofuku Y, Ueda T, Mase Y, Yokogawa M, Osawa M, Terashima Y, Matsushima K, Shimada I. NMR Analyses of the Interaction between CCR5 and Its Ligand Using Functional Reconstitution of CCR5 in Lipid Bilayers. *J Am Chem Soc.* 132:6768-77 2010.
535. Shono Y, Ueha S, Wang Y, Abe J, Kurachi M, Matsuno Y, Sugiyama T, Nagasawa T, Imamura M, Matsushima K. Bone marrow graft-versus-host disease: early destruction of hematopoietic niche following MHC-mismatched hematopoietic stem cell transplantation. *Blood.* 115:5401-11, 2010.
536. Esaki K, Terashima Y, Toda E, Yoshinaga S, Araki N, Matsushima K, Terasawa H. Expression and purification of human FROUNT, a common cytosolic regulator of CCR2 and CCR5. *Protein Expr Purif.* 77:86-91, 2010.
537. Otsuji M, Terashima Y, Ishihara S, Kuroda S, Matsushima K. A conceptual molecular network for chemotactic behaviors characterized by feedback of molecules cycling between the membrane and the cytosol. *Sci Signal.* 3:ra89. 2010.
538. Wada T, Sakai N, Sakai Y, Matsushima K, Kaneko S, Furuichi K. Involvement of bone-marrow-derived cells in kidney fibrosis. *Clin Exp Nephrol.* 15:8-13, 2011.
539. Iwata Y, Furuichi K, Kitagawa K, Hara A, Okumura T, Kokubo S, Shimizu K, Sakai N, Sagara A, Kurokawa Y, Ueha S, Matsushima K, Kaneko S, Wada T. Involvement of CD11b+ GR-1 low cells in autoimmune disorder in MRL-Fas lpr mouse. *Clin Exp Nephrol.* 14:411-7. 2010.
540. Hoshino A, Iimura T, Ueha S, Hanada S, Maruoka Y, Mayahara M, Suzuki K, Imai T, Ito M, Manome Y, Yasuhara M, Kirino T, Yamaguchi A, Matsushima K, Yamamoto K. Deficiency of chemokine receptor CCR1 causes osteopenia due to impaired functions of osteoclasts and osteoblasts. *J Biol Chem.* 285:28826-37. 2010.
541. Nakamoto Y, Mizukoshi E, Kitahara M, Arihara F, Sakai Y, Kakinoki K, Fujita Y, Marukawa Y, Arai K, Yamashita T, Mukaida N, Matsushima K, Matsui O, Kaneko S. Prolonged recurrence-free survival following OK432-stimulated dendritic cell transfer into hepatocellular carcinoma during transarterial embolization. *Clin Exp Immunol.* 163:165-77, 2011.
542. Ueha S, Shand FH, Matsushima K. Myeloid cell population dynamics in healthy and tumor-bearing mice. *Int Immunopharmacol.* 11:783-8, 2011.

543. Shimizu K, Furuichi K, Sakai N, Kitagawa K, Matsushima K, Mukaida N, Kaneko S, Wada T. Fractalkine and its receptor, CX3CR1, promote hypertensive interstitial fibrosis in the kidney. *Hypertens Res.* 34:747-52, 2011.
544. Toda M, Wang L, Ogura S, Torii M, Kurachi M, Kakimi K, Nishikawa H, Matsushima K, Shiku H, Kuribayashi K, Kato T. UV irradiation of immunized mice induces type 1 regulatory T cells that suppress tumor antigen specific cytotoxic T lymphocyte responses. *Int J Cancer.* 129:1126-36, 2011.
545. Kokubo S, Sakai N, Furuichi K, Toyama T, Kitajima S, Okumura T, Matsushima K, Kaneko S, Wada T. Activation of p38 mitogen-activated protein kinase promotes peritoneal fibrosis by regulating fibrocytes. *Perit Dial Int.* 32:10-9, 2011.
546. Kurachi M, Kurachi J, Suenaga F, Tsukui T, Abe J, Ueha S, Tomura M, Sugihara K, Takamura S, Kakimi K, Matsushima K. Chemokine receptor CXCR3 facilitates CD8+ T cell differentiation into short-lived effector cells leading to memory degeneration. *J Exp Med.* 208:1605-20, 2011.
547. Ogoshi K, Hashimoto S, Nakatani Y, Qu W, Oshima K, Tokunaga K, Sugano S, Hattori M, Morishita S, Matsushima K. Genome-wide profiling of DNA methylation in human cancer cells. *Genomics.* 98:280-7, 2011.
548. Kaneko K, Miyabe Y, Takayasu A, Fukuda S, Miyabe C, Ebisawa M, Yokoyama W, Watanabe K, Imai T, Muramoto K, Terashima Y, Sugihara T, Matsushima K, Miyasaka N, Nanki T. Chemerin activates fibroblast-like synoviocytes in patients with rheumatoid arthritis. *Arthritis Res Ther.* 13:R158, 2011.
549. Abe J, Ueha S, Yoneyama H, Shono Y, Kurachi M, Goto A, Fukayama M, Tomura M, Kakimi K, Matsushima K. B cells regulate antibody responses through the medullary remodeling of inflamed lymph nodes. *Int Immunol.* 24:17-27, 2012.
550. Ishida Y, Kimura A, Kuninaka Y, Inui M, Matsushima K, Mukaida N, Kondo T. Pivotal role of the CCL5/CCR5 interaction for recruitment of endothelial progenitor cells in mouse wound healing. *J Clin Invest.* 122:711-21, 2012.
551. Matsushima K, Nagai S. Unraveling the mystery of the hygiene hypothesis through *Helicobacter pylori* infection. *J Clin Invest.* 122:801-4, 2012.
552. Fujimura S, Takahashi H, Yuda K, Ueta T, Iriyama A, Inoue T, Kaburaki T, Tamaki Y, Matsushima K, Yanagi Y. Angiostatic effect of CXCR3 expressed on choroidal neovascularization. *Invest Ophthalmol Vis Sci.* 53:1999-2006, 2012.
553. Furuichi K, Shintani H, Sakai Y, Ochiya T, Matsushima K, Kaneko S, Wada T. Effects of adipose-derived mesenchymal cells on ischemia-reperfusion injury in kidney. *Clin Exp Nephrol.* 16:679-89, 2012.
554. Ueha S, Shand FH, Matsushima K. Cellular and molecular mechanisms of chronic inflammation-associated organ fibrosis. *Front Immunol.* 3:71, 2012.
555. Yamada Y, Okubo Y, Shimada A, Oikawa Y, Yamada S, Narumi S, Matsushima K, Itoh H. Acceleration of diabetes development in CXC chemokine receptor 3 (CXCR3)-deficient NOD mice. *Diabetologia.* 55:2238-45, 2012.

556. Furuichi K, Kokubo S, Hara A, Imamura R, Wang Q, Kitajima S, Toyama T, Okumura T, Matsushima K, Suda T, Mukaida N, Kaneko S, Wada T. Fas Ligand Has a Greater Impact than TNF- α on Apoptosis and Inflammation in Ischemic Acute Kidney Injury. *Nephron Extra.* 2:27-38, 2012.
557. Nagao K, Kobayashi T, Moro K, Ohyama M, Adachi T, Kitashima DY, Ueha S, Horiuchi K, Tanizaki H, Kabashima K, Kubo A, Cho YH, Clausen BE, Matsushima K, Suematsu M, Furtado GC, Lira SA, Farber JM, Udey MC, Amagai M. Stress-induced production of chemokines by hair follicles regulates the trafficking of dendritic cells in skin. *Nat Immunol.* 13:744-52, 2012.
558. Qu W, Hashimoto S, Shimada A, Nakatani Y, Ichikawa K, Saito TL, Ogoshi K, Matsushima K, Suzuki Y, Sugano S, Takeda H, Morishita S. Genome-wide genetic variations are highly correlated with proximal DNA methylation patterns. *Genome Res.* 22:1419-25, 2012.
559. Miyabe C, Miyabe Y, Miura NN, Takahashi K, Terashima Y, Toda E, Honda F, Morio T, Yamagata N, Ohno N, Shudo K, Suzuki J, Isobe M, Matsushima K, Tsuboi R, Miyasaka N, Nanki T. Am80, a retinoic acid receptor agonist, ameliorates murine vasculitis through the suppression of neutrophil migration and activation. *Arthritis Rheum.* 65:503-12, 2013.
560. Sekiguchi S, Kimura K, Chiyo T, Ohtsuki T, Tobita Y, Tokunaga Y, Yasui F, Tsukiyama-Kohara K, Wakita T, Tanaka T, Miyasaka M, Mizuno K, Hayashi Y, Hishima T, Matsushima K, Kohara M. Immunization with a recombinant vaccinia virus that encodes nonstructural proteins of the hepatitis C virus suppresses viral protein levels in mouse liver. *PLoS One.* 7:e51656, 2013.
561. Nakamura K, Doi K, Okamoto K, Arai S, Ueha S, Matsushima K, Nakajima S, Yahagi N, Noiri E. Specific antibody in IV immunoglobulin for postsplenectomy sepsis. *Crit Care Med.* 41(8):e163-70, 2013.
562. Hoshino A, Ueha S, Hanada S, Imai T, Ito M, Yamamoto K, Matsushima K, Yamaguchi A, Iimura T. Roles of chemokine receptor CX3CR1 in maintaining murine bone homeostasis through the regulation of both osteoblasts and osteoclasts. *J Cell Sci.* 126:1032-45, 2013.
563. Hashimoto S, Ogoshi K, Sasaki A, Abe J, Qu W, Nakatani Y, Ahsan B, Oshima K, Shand FH, Ametani A, Suzuki Y, Kaneko S, Wada T, Hattori M, Sugano S, Morishita S, Matsushima K. Coordinated changes in DNA methylation in antigen-specific memory CD4 T cells. *J Immunol.* 190:4076-91, 2013.
564. Hara A, Sakai N, Furuichi K, Sakai Y, Takeya M, Bucala R, Mukaida N, Takuwa Y, Matsushima K, Kaneko S, Wada T. CCL2/CCR2 augments the production of transforming growth factor-beta1, type 1 collagen and CCL2 by human CD45-/collagen 1-positive cells under high glucose concentrations. *Clin Exp Nephrol.* 17(6):793-804, 2013.
565. Onai N, Kurabayashi K, Hosoi-Amakie M, Toyama-Sorimachi N, Matsushima K, Inaba K, Ohteki T. A Clonogenic Progenitor with Prominent Plasmacytoid Dendritic Cell Developmental Potential. *Immunity.* 38:943-57, 2013.
566. Tsukui T, Ueha S, Abe J, Hashimoto S, Shichino S, Shimaoka T, Shand FH, Arakawa Y, Oshima K, Hattori M, Inagaki Y, Tomura M, Matsushima K. Qualitative rather than

- quantitative changes are hallmarks of fibroblasts in bleomycin-induced pulmonary fibrosis. *Am J Pathol.* 183:758-73, 2013.
567. Atarashi K, Tanoue T, Oshima K, Suda W, Nagano Y, Nishikawa H, Fukuda S, Saito T, Narushima S, Hase K, Kim S, Fritz JV, Wilmes P, Ueha S, Matsushima K, Ohno H, Olle B, Sakaguchi S, Taniguchi T, Morita H, Hattori M, Honda K. Treg induction by a rationally selected mixture of Clostridia strains from the human microbiota. *Nature.* 500:232-6, 2013.
568. Bachelierie F, Ben-Baruch A, Burkhardt AM, Combadiere C, Farber JM, Graham GJ, Horuk R, Sparre-Ulrich AH, Locati M, Luster AD, Mantovani A, Matsushima K, Murphy PM, Nibbs R, Nomiyama H, Power CA, Proudfoot AE, Rosenkilde MM, Rot A, Sozzani S, Thelen M, Yoshie O, Zlotnik A. International Union of Pharmacology. LXXXIX. Update on the Extended Family of Chemokine Receptors and Introducing a New Nomenclature for Atypical Chemokine Receptors. *Pharmacol Rev.* 2013, 66(1):1-79.
569. Wang SW, Wang SS, Wu DC, Lin YC, Ku CC, Wu CC, Chai CY, Lee JN, Tsai EM, Lin CL, Yang RC, Ko YC, Yu HS, Huo C, Chuu CP, Murayama Y, Nakamura Y, Hashimoto S, Matsushima K, Jin C, Eckner R, Lin CS, Saito S, Yokoyama KK. Androgen receptor-mediated apoptosis in bovine testicular induced pluripotent stem cells in response to phthalate esters. *Cell Death Dis.* 2013, 4:e907.
570. Hosoi A, Matsushita H, Shimizu K, Fujii SI, Ueha S, Abe J, Kurachi M, Maekawa R, Matsushima K, Kakimi K. Adoptive cytotoxic T lymphocyte therapy triggers a counter-regulatory immunosuppressive mechanism via recruitment of myeloid-derived suppressor cells. *Int J Cancer.* 34(8):1810-22, 2013.
571. Toda E, Terashima Y, Esaki K, Yoshinaga S, Sugihara M, Kofuku Y, Shimada I, Suwa M, Kanegasaki S, Terasawa H, Matsushima K. Identification of a binding element for the cytoplasmic regulator FROUNT in the membrane-proximal carboxy-terminal region of chemokine receptors CCR2 and CCR5. *Biochem J.* 2014, 457(2):313-22.
572. Ohyagi H, Onai N, Sato T, Yotsumoto S, Liu J, Akiba H, Yagita H, Atarashi K, Honda K, Roers A, Müller W, Kurabayashi K, Hosoi-Amaike M, Takahashi N, Hirokawa M, Matsushima K, Sawada K, Ohteki T. Monocyte-derived dendritic cells perform hemophagocytosis to fine-tune excessive immune responses. *Immunity.* 2013, 39(3):584-98.
573. Iwata Y, Furuichi K, Hashimoto S, Yokota K, Yasuda H, Sakai N, Kitajima S, Toyama T, Shinozaki Y, Sagara A, Matsushima K, Kaneko S, Wada T. Pro-inflammatory/Th1 gene expression shift in high glucose stimulated mesangial cells and tubular epithelial cells. *Biochem Biophys Res Commun.* 443(3):969-74, 2014.
574. Shono Y, Shiratori S, Kosugi-Kanaya M, Ueha S, Sugita J, Shigematsu A, Kondo T, Hashimoto D, Fujimoto K, Endo T, Nishio M, Hashino S, Matsuno Y, Matsushima K, Tanaka J, Imamura M, Teshima T. Bone marrow graft-versus-host disease: evaluation of its clinical impact on disrupted hematopoiesis after allogeneic hematopoietic stem cell transplantation. *Biol Blood Marrow Transplant.* 20(4):495-500, 2014.
575. Sasaki S, Baba T, Shinagawa K, Matsushima K, Mukaida N. Crucial involvement of the CCL3-CCR5 axis-mediated fibroblast accumulation in colitis-associated carcinogenesis in mice. *Int J Cancer.* 135(6):1297-306, 2014.

576. Lin YC, Kuo KK, Wuputra K, Lin SH, Ku CC, Yang YH, Wang SW, Wang SW, Wu DC, Wu CC, Chai CY, Lin CL, Lin CS, Kajitani M, Miyoshi H, Nakamura Y, Hashimoto S, Matsushima K, Jin C, Huang SK, Saito S, Yokoyama KK. Bovine induced pluripotent stem cells are more resistant to apoptosis than testicular cells in response to mono-(2-ethylhexyl) phthalate. *Int J Mol Sci.* 15(3):5011-31, 2014.
577. Sumida H, Yanagida K, Kita Y, Abe J, Matsushima K, Nakamura M, Ishii S, Sato S, Shimizu T. Interplay between CXCR2 and BLT1 facilitates neutrophil infiltration and resultant keratinocyte activation in a murine model of imiquimod-induced psoriasis. *J Immunol.* 192(9):4361-9, 2014.
578. Yasui F, Kohara M, Kitabatake M, Nishiwaki T, Fujii H, Tateno C, Yoneda M, Morita K, Matsushima K, Koyasu S, Kai C. Phagocytic cells contribute to the antibody-mediated elimination of pulmonary-infected SARS coronavirus. *Virology.* 454-455:157-68, 2014.
579. Shand FH, Ueha S, Otsuji M, Koid SS, Shichino S, Tsukui T, Kosugi-Kanaya M, Abe J, Tomura M, Ziogas J, Matsushima K. Tracking of intertissue migration reveals the origins of tumor-infiltrating monocytes. *Proc Natl Acad Sci U S A.* 111(21):7771-6, 2014.
580. Kitahara M, Mizukoshi E, Nakamoto Y, Mukaida N, Matsushima K, Kaneko S. Efficient generation of highly immunocompetent dendritic cells from peripheral blood of patients with hepatitis C virus-related hepatocellular carcinoma. *Int Immunopharmacol.* 21(2):346-53, 2014.
581. Nakamura R, Tsukahara T, Qu W, Ichikawa K, Otsuka T, Ogoshi K, Saito TL, Matsushima K, Sugano S, Hashimoto S, Suzuki Y, Morishita S, Takeda H. Large hypomethylated domains serve as strong repressive machinery for key developmental genes in vertebrates. *Development.* 141(13):2568-80, 2014.
582. Kanegasaki S, Matsushima K, Shiraishi K, Nakagawa K, Tsuchiya T. Macrophage Inflammatory Protein Derivative ECI301 Enhances the Alarmin-Associated Abscopal Benefits of Tumor Radiotherapy. *Cancer Res.* 15;74(18):5070-8, 2014.
583. Abe J, Shichino S, Ueha S, Hashimoto S, Tomura M, Inagaki Y, Stein JV, Matsushima K. Lymph node stromal cells negatively regulate antigen-specific CD4+ T cell responses. *J Immunol.* 193(4):1636-44, 2014.
584. Matsushita H, Hosoi A, Ueha S, Abe J, Fujieda N, Tomura M, Maekawa R, Matsushima K, Ohara O, Kakimi K. Cytotoxic T lymphocytes block tumor growth both by lytic activity and IFN- γ -dependent cell cycle arrest. *Cancer Immunol Res.* 3:26-36, 2014.
585. Tomura M, Hata A, Matsuoka S, Shand FH, Nakanishi Y, Ikebuchi R, Ueha S, Tsutsui H, Inaba K, Matsushima K, Miyawaki A, Kabashima K, Watanabe T, Kanagawa O. Tracking and quantification of dendritic cell migration and antigen trafficking between the skin and lymph nodes. *Sci Rep.* 4:6030, 2014.
586. Yoshie O, Matsushima K. CCR4 and its ligands: from bench to bedside. *Int Immunol.* 27:11-20, 2014.
587. Esaki K, Yoshinaga S, Tsuji T, Toda E, Terashima Y, Saitoh T, Kohda D, Kohno T, Osawa M, Ueda T, Shimada I, Matsushima K, Terasawa H. Structural basis for the binding of the

- membrane-proximal C-terminal region of chemokine receptor CCR2 with the cytosolic regulator FROUNT. *FEBS J.* 281(24):5552-66, 2014.
588. Suenaga F, Ueha S, Abe J, Kosugi-Kanaya M, Wang Y, Yokoyama A, Shono Y, Shand FH, Morishita Y, Kunisawa J, Sato S, Kiyono H, Matsushima K. Loss of Lymph Node Fibroblastic Reticular Cells and High Endothelial Cells Is Associated with Humoral Immunodeficiency in Mouse Graft-versus-Host Disease. *J Immunol.* 194(1):398-406, 2015.
589. Ueha S, Yokochi S, Ishiwata Y, Ogiwara H, Chand K, Nakajima T, Hachiga K, Shichino S, Terashima Y, Toda E, Shand FH, Kakimi K, Ito S, Matsushima K. Robust Antitumor Effects of Combined Anti-CD4-Depleting Antibody and Anti-PD-1/PD-L1 Immune Checkpoint Antibody Treatment in Mice. *Cancer Immunol Res.* 3(6):631-40, 2015.
590. Hirano K, Hosoi A, Matsushita H, Ino T, Ueha S, Matsushima K, Seto Y, Kakimi K. The nitric oxide radical scavenger carboxy-PTIO reduces the immunosuppressive activity of myeloid-derived suppressor cells and potentiates the antitumor activity of adoptive cytotoxic T lymphocyte immunotherapy. *Oncoimmunology.* 4(8):e1019195, 2015.
591. Shichino S, Abe J, Ueha S, Otsuji M, Tsukui T, Kosugi-Kanaya M, Shand FH, Hashimoto S, Suzuki HI, Morikawa T, Inagaki Y, Matsushima K. Reduced supply of monocyte-derived macrophages leads to a transition from nodular to diffuse lesions and tissue cell activation in silica-induced pulmonary fibrosis in mice. *Am J Pathol.* 185(11):2923-38, 2015.
592. Tsukui T, Ueha S, Shichino S, Inagaki Y, Matsushima K. Intratracheal cell transfer demonstrates the profibrotic potential of resident fibroblasts in pulmonary fibrosis. *Am J Pathol.* 185(11):2939-48, 2015.
593. Ueha R, Ueha S, Kondo K, Sakamoto T, Kikuta S, Kanaya K, Nishijima H, Matsushima K, Yamasoba T. Damage to Olfactory Progenitor Cells Is Involved in Cigarette Smoke-Induced Olfactory Dysfunction in Mice. *Am J Pathol.* 186(3):579-86, 2016.
594. Moro T, Nakao S, Sumiyoshi H, Ishii T, Miyazawa M, Ishii N, Sato T, Iida Y, Okada Y, Tanaka M, Hayashi H, Ueha S, Matsushima K, Inagaki Y. A Combination of Mitochondrial Oxidative Stress and Excess Fat/Calorie Intake Accelerates Steatohepatitis by Enhancing Hepatic CC Chemokine Production in Mice. *PLoS One.* 11(1):e0146592, 2016.
595. Park AM, Kanai K, Itoh T, Sato T, Tsukui T, Inagaki Y, Selman M, Matsushima K, Yoshie O. Heat Shock Protein 27 Plays a Pivotal Role in Myofibroblast Differentiation and in the Development of Bleomycin-Induced Pulmonary Fibrosis. *PLoS One.* 11(2):e0148998, 2016.
596. Matsuo Y, Mizoguchi F, Saito T, Kawahata K, Ueha S, Matsushima K, Inagaki Y, Miyasaka N, Kohsaka H. Local fibroblast proliferation but not influx is responsible for synovial hyperplasia in a murine model of rheumatoid arthritis. *Biochem Biophys Res Commun.* 470(3):504-9, 2016.
597. Yasui F, Kohara M, Kitabatake M, Nishiwaki T, Fujii H, Tateno C, Yoneda M, Morita K, Matsushima K, Koyasu S, Kai C. Corrigendum to 'Phagocytic cells contribute to the antibody-mediated elimination of pulmonary-infected SARS coronavirus' [*Virology* (2014) 157-168]. *Virology.* 499:397-398, 2016.
598. Tan X, Fujii K, Manabe I, Nishida J, Yamagishi R, Terashima Y, Matsushima K, Kaburaki T, Nagai R, Yanagi Y. Choroidal Neovascularization Is Inhibited in Splenic-Denervated or

- Splenectomized Mice with a Concomitant Decrease in Intraocular Macrophage. *PLoS One*. 11(8):e0160985, 2016.
599. Ueha R, Ueha S, Sakamoto T, Kanaya K, Suzukawa K, Nishijima H, Kikuta S, Kondo K, Matsushima K, Yamasoba T. Cigarette Smoke Delays Regeneration of the Olfactory Epithelium in Mice. *Neurotox Res*. 30(2):213-24, 2016.
600. Ueha R, Ueha S, Kondo K, Nito T, Fujimaki Y, Nishijima H, Tsunoda K, Shand FH, Matsushima K, Yamasoba T. Laryngeal mucus hypersecretion is exacerbated after smoking cessation and ameliorated by glucocorticoid administration. *Toxicol Lett*. 265:140-146, 2017.
601. Fujino M, Sato H, Okamura T, Uda A, Takeda S, Ahmed N, Shichino S, Shiino T, Saito Y, Watanabe S, Sugimoto C, Kuroda M, Ato M, Nagai Y, Izumo S, Matsushima K, Miyazawa M, Ansari AA, Villinger F, Mori K. Simian Immunodeficiency Virus Targeting of CXCR3⁺ CD4⁺ T Cells in Secondary Lymphoid Organs Is Associated with Robust CXCL10 Expression in Monocyte/Macrophage Subsets. *J Virol*. 91(13). pii: e00439-17, 2017.
602. Sonoda A, Yoshinaga S, Yunoki K, Ezaki S, Yano K, Takeda M, Toda E, Terashima Y, Matsushima K, Terasawa H. Identification and Preparation of a Novel Chemokine Receptor-Binding Domain in the Cytoplasmic Regulator FROUNT. *Mol Biotechnol*. 59(4-5):141-150, 2017.
603. Ueha S, Yokochi S, Ishiwata Y, Kosugi-Kanaya M, Shono Y, Shibayama S, Ito S, Matsushima K. Combination of anti-CD4 antibody treatment and donor lymphocyte infusion ameliorates graft-versus-host disease while preserving graft-versus-tumor effects in murine allogeneic hematopoietic stem cell transplantation. *Cancer Sci*. 108(10):1967-1973, 2017.
604. Hashimoto S, Tabuchi Y, Yurino H, Hirohashi Y, Deshimaru S, Asano T, Mariya T, Oshima K, Takamura Y, Ukita Y, Ametani A, Kondo N, Monma N, Takeda T, Misu S, Okayama T, Ikeo K, Saito T, Kaneko S, Suzuki Y, Hattori M, Matsushima K, Torigoe T. Comprehensive single-cell transcriptome analysis reveals heterogeneity in endometrioid adenocarcinoma tissues. *Sci Rep*. 7(1):14225, 2017.
605. Lee JW, Hoshino A, Inoue K, Saitou T, Uehara S, Kobayashi Y, Ueha S, Matsushima K, Yamaguchi A, Imai Y, Iimura T. The HIV co-receptor CCR5 regulates osteoclast function. *Nat Commun*. 8(1):2226, 2017.
606. Kosugi-Kanaya M, Ueha S, Abe J, Shichino S, Shand FHW, Morikawa T, Kurachi M, Shono Y, Sudo N, Yamashita A, Suenaga F, Yokoyama A, Yong W, Imamura M, Teshima T, Matsushima K. Long-Lasting Graft-Derived Donor T Cells Contribute to the Pathogenesis of Chronic Graft-versus-Host Disease in Mice. *Front Immunol*. 8:1842, 2017.
607. Yunoki K, Yoshinaga S, Takeda M, Nagano R, Tsuchiya Y, Sonoda A, Tsuji T, Hirakane M, Toda E, Terashima Y, Matsushima K, Terasawa H. Efficient identification of compounds suppressing protein precipitation via solvent screening using serial deletion mutants of the target protein. *Genes Cells*. 23(2):70-79, 2018.
608. Hosoi A, Takeda K, Nagaoka K, Iino T, Matsushita H, Ueha S, Aoki S, Matsushima K, Kubo M, Morikawa T, Kitaura K, Suzuki R, Kakimi K. Increased diversity with reduced "diversity evenness" of tumor infiltrating T-cells for the successful cancer immunotherapy. *Sci Rep*. 8(1):1058, 2018.

609. Ueta H, Kitazawa Y, Sawanobori Y, Ueno T, Ueha S, Matsushima K, Matsuno K. Single blood transfusion induces the production of donor-specific alloantibodies and regulatory T cells mainly in the spleen. *Int Immunol.* 30(2):53-67, 2018.
610. Yoshinaga S, Ishida N, Tsuji T, Sonoda A, Yunoki K, Takeda M, Toda E, Terashima Y, Matsushima K, Terasawa H. ¹H, ¹³C and ¹⁵N resonance assignments for a chemokine receptor-binding domain of FROUNT, a cytoplasmic regulator of chemotaxis. *Biomol NMR Assign.* 12(2):259-262, 2018.
611. Ueha R, Shichino S, Ueha S, Kondo K, Kikuta S, Nishijima H, Matsushima K, Yamasoba T. Reduction of Proliferating Olfactory Cells and Low Expression of Extracellular Matrix Genes Are Hallmarks of the Aged Olfactory Mucosa. *Front Aging Neurosci.* 10:86, 2018.
612. Souma K, Shichino S, Hashimoto S, Ueha S, Tsukui T, Nakajima T, Suzuki HI, Shand FHW, Inagaki Y, Nagase T, Matsushima K. Lung fibroblasts express a miR-19a-19b-20a sub-cluster to suppress TGF- β -associated fibroblast activation in murine pulmonary fibrosis. *Sci Rep.* 8(1):16642, 2018.
613. Shichino S, Ueha S, Hashimoto S, Otsuji M, Abe J, Tsukui T, Deshimaru S, Nakajima T, Kosugi-Kanaya M, Shand FH, Inagaki Y, Shimano H, Matsushima K. Transcriptome network analysis identifies protective role of the LXR/SREBP-1c axis in murine pulmonary fibrosis. *JCI Insight.* 4(1), 2019.
614. Shiraishi K, Shichino S, Ueha S, Nakajima T, Hashimoto S, Yamazaki S, Matsushima K. Mesenchymal-Epithelial Interactome Analysis Reveals Essential Factors Required for Fibroblast-Free Alveolosphere Formation. *iScience.* 11:318-333, 2019.
615. Chen CY, Ueha S, Ishiwata Y, Yokochi S, Yang D, Oppenheim JJ, Ogiwara H, Shichino S, Deshimaru S, Shand FH, Shibayama S, Matsushima K. Combined treatment with HMGN1 and anti-CD4 depleting antibody reverses T cell exhaustion and exerts robust anti-tumor effects in mice. *J Immunother.* 7(1):21, 2019.
616. Aoki H, Ueha S, Shichino S, Ogiwara H, Hashimoto S, Kakimi K, Ito S, Matsushima K. TCR repertoire analysis reveals mobilization of novel CD8⁺ T cell clones into the Cancer-Immunity Cycle following anti-CD4 antibody administration. *Front Immunol.* 9:3185, 2019.
617. Shiraishi K, Shichino S, Tsukui T, Hashimoto S, Ueha S, Matsushima K. Engraftment and proliferation potential of embryonic lung tissue cells in irradiated mice with emphysema. *Sci Rep.* 9(1):3657, 2019.
618. Tsukui T, Ueha S, Shichino S, Hashimoto S, Nakajima T, Shiraishi K, Kihara M, Kiyonari H, Inagaki Y, Matsushima K. Gli signaling pathway modulates fibroblast activation and facilitates scar formation in pulmonary fibrosis. *Biochem Biophys Res Commun.* 514(3):684-690, 2019.
619. Shiraishi K, Nakajima T, Shichino S, Deshimaru S, Matsushima K, Ueha S. In vitro expansion of endogenous human alveolar epithelial type II cells in fibroblast-free spheroid culture. *Biochem Biophys Res Commun.* 515(4):579-585, 2019.

620. Kitazawa Y, Ueta H, Sawanobori Y, Katakai T, Yoneyama H, Ueha S, Matsushima K, Tokuda N, Matsuno K. Novel Targeting to XCR1⁺ Dendritic Cells Using Allogeneic T Cells for Polytopical Antibody Responses in the Lymph Nodes. *Front Immunol.* 10:1195, 2019.
621. Shitara K, Ueha S, Shichino S, Aoki H, Ogiwara H, Nakatsura T, Suzuki T, Shimomura M, Yoshikawa T, Shoda K, Kitano S, Yamashita M, Nakayama T, Sato A, Kuroda S, Wakabayashi M, Nomura S, Yokochi S, Ito S, Matsushima K, Doi T. First-in-human phase 1 study of IT1208, a defucosylated humanized anti-CD4 depleting antibody, in patients with advanced solid tumors. *J Immunother Cancer.* 7(1):195, 2019.
622. Wein AN, McMaster SR, Takamura S, Dunbar PR, Cartwright EK, Hayward SL, McManus DT, Shimaoka T, Ueha S, Tsukui T, Masumoto T, Kurachi M, Matsushima K, Kohlmeier JE. CXCR6 regulates localization of tissue-resident memory CD8 T cells to the airways. *J Exp Med.* 216(12):2748-2762, 2019.
623. Takamura S, Kato S, Motozono C, Shimaoka T, Ueha S, Matsuo K, Miyauchi K, Masumoto T, Katsushima A, Nakayama T, Tomura M, Matsushima K, Kubo M, Miyazawa M. Interstitial-resident memory CD8⁺ T cells sustain frontline epithelial memory in the lung. *J Exp Med.* 216(12):2736-2747, 2019.
624. Iwata Y, Satou K, Furuichi K, Yoneda I, Matsumura T, Yutani M, Fujinaga Y, Hase A, Morita H, Ohta T, Senda Y, Sakai-Takemori Y, Wada T, Fujita S, Miyake T, Yasuda H, Sakai N, Kitajima S, Toyama T, Shinozaki Y, Sagara A, Miyagawa T, Hara A, Shimizu M, Kamikawa Y, Ikeo K, Shichino S, Ueha S, Nakajima T, Matsushima K, Kaneko S, Wada T. Collagen adhesion gene is associated with bloodstream infections caused by methicillin-resistant *Staphylococcus aureus*. *Int J Infect Dis.* 91:22-31, 2020.
625. Terashima Y, Toda E, Itakura M, Otsuji M, Yoshinaga S, Okumura K, Shand FHW, Komohara Y, Takeda M, Kokubo K, Chen MC, Yokoi S, Rokutan H, Kofuku Y, Ohnishi K, Ohira M, Iizasa T, Nakano H, Okabe T, Kojima H, Shimizu A, Kanegasaki S, Zhang MR, Shimada I, Nagase H, Terasawa H, Matsushima K. Targeting FROUNT with disulfiram suppresses macrophage accumulation and its tumor-promoting properties. *Nat Commun.* 11(1):609, 2020.
626. Ishii H, Yamasaki T, Yui J, Zhang Y, Hanyu M, Ogawa M, Nengaki N, Tsuji AB, Terashima Y, Matsushima K, Zhang MR. Radiosynthesis of [thiocarbonyl-¹¹C]disulfiram and its first PET study in mice. *Bioorg Med Chem Lett.* 30(6):126998, 2020.
627. Iwata Y, Sakai N, Yoneda I, Satou K, Furuichi K, Senda Y, Sakai-Takemori Y, Wada T, Fujita S, Ogura H, Sato K, Minami T, Yamaguchi K, Kitajima S, Toyama T, Yamamura Y, Miyagawa T, Hara A, Shimizu M, Sakai Y, Ikeo K, Shichino S, Ueha S, Nakajima T, Matsushima K, Wada T. The increased frequency of methicillin-resistant *Staphylococcus aureus* with low MIC of beta-lactam antibiotics isolated from hospitalized patients. *J Infect Chemother.* 26(6):604-610, 2020.
628. Watanabe S, Fujino M, Saito Y, Ahmed N, Sato H, Sugimoto C, Okamura T, Hanaki K, Nakayama EE, Shioda T, Matsushima K, Ansari AA, Villinger F, Mori K. Protective Immune Responses Elicited by Deglycosylated Live-Attenuated Simian Immunodeficiency Virus Vaccine Are Associated with IL-15 Effector Functions. *J Immunol.* 205(5):1331-1344, 2020.

629. Mizoguchi I, Ohashi M, Hasegawa H, Chiba Y, Orii N, Inoue S, Kawana C, Xu M, Sudo K, Fujita K, Kuroda M, Hashimoto SI, Matsushima K, Yoshimoto T. EBV-induced gene 3 augments IL-23R α protein expression through a chaperone calnexin. *J Clin Invest.* 122732, 2020.
630. Tsuji T, Yoshinaga S, Takeda M, Sato T, Sonoda A, Ishida N, Yunoki K, Toda E, Terashima Y, Matsushima K, Terasawa H. Rational Design of Monodispersed Mutants of Proteins by Identifying Aggregation Contact Sites Using Solubilizing Agents. *Biochemistry.* 59(39):3639-3649, 2020.
631. Omori S, Wang TW, Johmura Y, Kanai T, Nakano Y, Kido T, Susaki EA, Nakajima T, Shichino S, Ueha S, Ozawa M, Yokote K, Kumamoto S, Nishiyama A, Sakamoto T, Yamaguchi K, Hatakeyama S, Shimizu E, Katayama K, Yamada Y, Yamazaki S, Iwasaki K, Miyoshi C, Funato H, Yanagisawa M, Ueno H, Imoto S, Furukawa Y, Yoshida N, Matsushima K, Ueda HR, Miyajima A, Nakanishi M. Generation of a p16 Reporter Mouse and Its Use to Characterize and Target p16^{high} Cells In Vivo. *Cell Metab.* S1550-4131(20)30484-8, 2020.
632. Ueta H, Xu XD, Yu B, Kitazawa Y, Yu E, Hara Y, Morita-Nakagawa M, Zhou S, Sawanobori Y, Ueha S, Rokutan K, Tanaka T, Tokuda N, Matsushima K, Matsuno K. Suppression of liver transplant rejection by anti-donor MHC antibodies via \square depletion of donor immunogenic dendritic cells. *Int Immunol.* dxaa076, 2020.
633. Hashimoto S, Shirasaki T, Yamashita T, Iwabuchi S, Suzuki Y, Takamura Y, Ukita Y, Deshimaru S, Okayama T, Ieko K, Kuroki K, Kawaguchi K, Mizukoshi E, Matsushima K, Honda M, Kaneko S. DOCK11 and DENND2A play pivotal roles in the maintenance of hepatitis B virus in host cells. *PLoS One.* 16(2):e0246313, 2021.
634. Iwata Y, Sakai N, Yoneda I, Senda Y, Sakai-Takemori Y, Oshima M, Nakagawa-Yoneda S, Ogura H, Sato K, Minami T, Kitajima S, Toyama T, Yamamura Y, Miyagawa T, Hara A, Shimizu M, Furuichi K, Matsushima K, Wada T. D-Serine inhibits the attachment and biofilm formation of methicillin-resistant *Staphylococcus aureus*. *Biochem Biophys Res Commun.* 537:50-56, 2021.
635. Aoki H, Ueha S, Shichino S, Ogiwara H, Shitara K, Shimomura M, Suzuki T, Nakatsura T, Yamashita M, Kitano S, Kuroda S, Wakabayashi M, Kurachi M, Ito S, Doi T, Matsushima K. Transient depletion of CD4⁺ cells induces drastic remodeling of the repertoire of T-cell receptors in gastrointestinal cancer. *Cancer Immunol Res.* 9:624-636, 2021.
636. Chen CY, Ueha S, Ishiwata Y, Shichino S, Yokochi S, Yang D, Oppenheim JJ, Ogiwara H, Deshimaru S, Kanno Y, Aoki H, Ogawa T, Shibayama S, Kouji Matsushima. Combining an alarmin HMG1 peptide with PD-L1 blockade results in robust antitumor effects with a concomitant increase of stem-like/progenitor exhausted CD8⁺ T cells. *Cancer Immunol Res.* 9:1214-1228, 2021.
637. Aoki H, Ueha S, Nakamura Y, Shichino S, Nakajima H, Shimomura M, Sato A, Nakatsura T, Yoshino T, Kouji Matsushima. Greater extent of blood-tumor TCR repertoire overlap is associated with favorable clinical responses to PD-1 blockade. *Cancer Sci.* 112:2993-3004, 2021.

638. Ogawa T, Shichino S, Ueha S, Kouji Matsushima. Macrophages in lung fibrosis. *Int Immunol.* 33:665-671, 2021.